

U.S. Army Public Health Command (Provisional)

INJURY PREVENTION REPORT NO. 12-HF-0893-10
INJURY INCIDENCE AND INJURY RISK FACTORS AMONG
SOLDIERS IN THE U.S. ARMY ORDNANCE SCHOOL
ABERDEEN PROVING GROUND, MARYLAND
JANUARY 2000–JUNE 2003

Approved for public release; distribution unlimited.

Preventive Medicine Survey: 40-5f1

U
S
A
P
H
C

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>				
1. REPORT DATE (DD-MM-YYYY) July 2009		2. REPORT TYPE Final		3. DATES COVERED (From – To) January 2000-June 2003
4. TITLE AND SUBTITLE Injury Incidence and Injury Risk Factors Among Soldiers in the United States Army Ordnance School			5a. CONTRACT NUMBER	
			5b. GRANT NUMBER	
			5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Tyson Grier, Stephanie Morrison, Joseph J Knapik, Michelle Canham-Chervak, Bruce H Jones			5d. PROJECT NUMBER	
			5e. TASK NUMBER	
			5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Center for Health Promotion and Preventive Medicine Aberdeen Proving Ground, MD			8. PERFORMING ORGANIZATION REPORT NUMBER 12-HF-0893-09	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)	
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited				
13. SUPPLEMENTARY NOTES				
14. ABSTRACT <p>This study examined risk factors for time-loss injuries during U.S. Army Ordnance School Advanced Individual Training (AIT). Participants were soldiers (n= 3757 men, n=498 women) attending Ordnance AIT from January 2000 to June 2003. Injuries were obtained from an injury surveillance system in the medical clinic serving the AIT soldiers. Potential injury risk factors included entry-level fitness, demographics, and lifestyle variables. Fitness variables included maximal effort performance on push-ups, sit-ups and a 2-mile run. A health questionnaire provided data on age, race, rank, current self reported injury, current self reported illness, and tobacco use. Fitness variables were converted to four quartiles (Q) based on the distribution of scores (Q1=high performance to Q4=low performance). AIT entry and completion dates were obtained from an Army personnel system. Backward stepping Cox regression examined associations between time-loss injury and potential injury risk factors. Multivariate hazard ratios (MHR) and 95% confidence intervals (95%CI) were calculated. Cumulative time-loss injury incidence was 31% for men and 54% for women. For men, higher risk of injury resulting in time-loss was independently associated with race (MHR (Native American/Caucasian) = 1.4, 95%CI = 1.1-1.7), a current self reported injury (MHR (yes/no) = 2.2, 95%CI = 1.8-2.7), smoking before entering the Army (MHR (≤ 10 cigarettes/nonsmokers) = 1.3, 95%CI = 1.0-1.5, MHR (10-20 cigarettes/nonsmokers) = 1.5, 95%CI = 1.2-1.7, MHR (> 20 cigarettes/nonsmokers) = 1.9, 95%CI = 1.6-2.2), lower sit-up performance (MHR (Q4/Q1) = 1.2, 95%CI = 1.0-1.5), and slower 2-mile run times (MHR (Q4/Q1) = 1.4, 95%CI = 1.2-1.7). For women, higher risk of injury was independently associated with race (MHR (Black/Caucasian) = 0.7, 95%CI = 0.5-0.9), a current self reported injury (MHR (yes/no) = 1.6, 95%CI = 1.1-2.3), and slower 2-mile run times (MHR (Q4/Q1) = 2.2, 95%CI = 1.5-3.1). Risk factors for time-loss injury in Ordnance AIT for both men and women included race, current self reported injuries, and lower aerobic fitness. Smoking cessation classes and fitness training prior to entry are potential strategies to reduce injuries among soldiers in Ordnance School AIT.</p>				
15. SUBJECT TERMS Injury, smokeless tobacco, tobacco, military, ordnance school				
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified		
				19a. NAME OF RESPONSIBLE PERSON Tyson Grier
				19b. TELEPHONE NUMBER (include area code) 410-436-5450/3534



DEPARTMENT OF THE ARMY
US ARMY PUBLIC HEALTH COMMAND (PROVISIONAL)
5158 BLACKHAWK ROAD
ABERDEEN PROVING GROUND MD 21010-5403

MCHB-TS-DI

EXECUTIVE SUMMARY
INJURY PREVENTION REPORT NO. 12-HF-0893-10
INJURY INCIDENCE AND INJURY RISK FACTORS AMONG SOLDIERS IN THE
U.S. ARMY ORDNANCE SCHOOL
ABERDEEN PROVING GROUND, MARYLAND
JANUARY 2000–JUNE 2003

1. PURPOSE. From January 2000 to June 2003, the U.S. Army Public Health Command (Provisional), formerly known as the U.S. Army Center for Health Promotion and Preventive Medicine, and the Kirk U.S. Army Health Clinic collaborated with the 143rd Ordnance Battalion at Aberdeen Proving Ground (APG), Maryland to identify injuries and injury risk factors during advanced individual training (AIT). This report provides the results of this collaboration by addressing injury rates and injury risk factors among Soldiers in AIT at the Ordnance School.

2. METHODS.

a. Participants were Army Service members attending AIT at the Edgewood Area of APG from January 2000 to June 2003. These Service members were training to qualify for one of five different Military Occupational Specialties (MOSs). These MOSs included track vehicle repairer (MOS 63H), wheeled vehicle repairer (MOS 63W), self-propelled field artillery system mechanic (MOS 63D), fuel and electrical system repairer (MOS 63G), and track vehicle mechanic (MOS 63Y).

b. On arrival at APG, each Service member was asked to complete a Soldier health in-processing questionnaire. The questionnaire included questions about the demographics and lifestyle characteristics of Service members. Information requested included whether or not the student currently had an injury or illness perceived to affect their AIT performance, history of tobacco use, date of birth, gender, rank, race, and basic combat training (BCT) location. Army Physical Fitness Test (APFT) data were obtained from the 143rd Ordnance Battalion Training and Operations (S-3) office.

c. Injuries occurring during training were obtained from an injury surveillance system in the medical clinic serving the AIT Soldiers. Every time a Soldier reported to the medical clinic, a medical provider would fill out an injury sheet. The injury sheet had boxes for the medical provider to check that indicated the type of injury that had occurred and the number of profile or quarter days given to the Soldier. Injuries from the data sheet were classified into four categories: time-loss injuries, overuse time-loss injuries, lower extremity overuse time-loss injuries and traumatic time-loss injuries. Time-loss injuries include an injury of any type plus a profile of one or more days.

d. The Statistical Package for the Social Sciences, Version 16.0, was used for statistical analysis. Potential risk factors for time-loss injuries, overuse time-loss injuries, lower extremity overuse time-loss injuries and traumatic time-loss injuries were explored using Cox regression (separate models were developed for each injury category). Hazard ratios and 95 percent confidence intervals (CIs) were calculated for each potential injury risk factor. Variables from the univariate analysis with a statistical significance of $p \leq 0.10$ were selected for a backward-stepping multivariate Cox regression. Multivariate hazard ratios (MHRs) and 95 percent CIs were calculated. For the Cox regression, APFT scores were converted to four quartiles (Qs) based on the distribution of scores where Q1 = high performance and Q4=low performance.

3. RESULTS.

a. There were 3757 men and 498 women involved with the project. A majority of the subjects were Caucasian men between the ages of 17 and 19 who had attended BCT at Fort Knox and were wheeled vehicle repairers (MOS 63W) of lower military rank (E1) and nonsmokers and non-smokeless tobacco users. During the course of their Ordnance School training, 31 percent of men and 54 percent of women had one or more time-loss injuries; the time-loss injury rate for men over this time period was 34.9 injuries/10,000 person-days, and the rate for women was 60.8 injuries/10,000 person-days.

b. Multivariate logistic regression results follow.

(1) Time-Loss Injury.

(a) For men, a higher risk of time-loss injury was associated with the Native American race (MHR (Native American/Caucasian) = 1.4, 95 percent CI = 1.1–1.7); a current self-reported injury (MHR (yes/no) = 2.2, 95 percent CI = 1.8–2.7); smoking before entering the Army (MHR (≤ 10 cigarettes/nonsmokers) = 1.3, 95 percent CI = 1.0–1.5, MHR (10-20 cigarettes/nonsmokers) = 1.5, 95 percent CI = 1.2–1.7, MHR (> 20 cigarettes/nonsmokers) = 1.9, 95 percent CI = 1.6–2.2); lower sit-up performance (MHR (Q4/Q1) = 1.2, 95 percent CI = 1.0–1.5); and slower 2-mile run times (MHR (Q4/Q1) = 1.4, 95 percent CI = 1.2–1.7).

(b) For women, a higher risk of time-loss injury was associated with Caucasian race (MHR (Black/Caucasian) = 0.7, 95 percent CI = 0.5–0.9); a current self-reported injury (MHR (yes/no) = 1.6, 95 percent CI = 1.1–2.3); and slower 2-mile run times (MHR (Q4/Q1) = 2.2, 95 percent CI = 1.5–3.1).

(2) Overuse Time-Loss Injury.

(a) For men, a higher risk of overuse time-loss injury was associated with MOS (MHR (fuel and electrical system repairer (MOS 63G)/track vehicle repairer (MOS 63H)) = 1.5, 95 percent CI = 1.1–2.1, MHR (track vehicle mechanic (MOS 63Y)/track vehicle repairer (MOS 63H)) = 1.4, 95 percent CI = 1.1–1.8); a current self-reported injury (MHR (yes/no) = 2.2, 95 percent CI = 1.8–2.8); smoking before entering the Army (MHR (10–20 cigarettes/nonsmokers) = 1.7, 95 percent CI = 1.4–2.0, MHR (>20 cigarettes/nonsmokers) = 1.9, 95 percent CI = 1.6–2.4); lower sit-up performance (MHR (Q4/Q1) = 1.4, 95 percent CI = 1.1–1.7); and slower 2-mile run times (MHR (Q4/Q1) = 1.5, 95 percent CI = 1.2–1.8).

(b) For women, a higher risk of overuse time-loss injury was independently associated with age (MHR (20–24/17–19) = 0.73, 95 percent CI = 0.5–1.00); Caucasian race (MHR (Black/Caucasians) = 0.6, 95 percent CI = 0.4–0.9); attending basic training at Fort Leonard Wood (MHR (Fort Leonard Wood/Fort Jackson) = 1.6, 95 percent CI = 1.1–2.3); a current self-reported injury (MHR (yes/no) = 1.5, 95 percent CI = 1.0–2.1); and slower 2-mile run times (MHR (Q4/Q1) = 2.0, 95 percent CI = 1.3–2.9).

(3) Lower Extremity Overuse Time-Loss Injury.

(a) For men, a higher risk of time-losslower extremity overuse time-loss injury was associated with a current self-reported injury (MHR (yes/no) = 2.0, 95 percent CI = 1.5–2.5); smoking before entering the Army (MHR (10–20 cigarettes/nonsmokers) = 1.7, 95 percent CI = 1.4–2.1, MHR (>20 cigarettes/nonsmokers) = 2.1, 95 percent CI = 1.7–2.6); lower sit-up performance (MHR (Q4/Q1) = 1.5, 95 percent CI = 1.1–1.9); and slower 2-mile run times (MHR (Q4/Q1) = 1.5, 95 percent CI = 1.2–1.9).

(b) For women, a higher risk of time-losslower extremity overuse time-loss injury was associated with attending basic training at Fort Leonard Wood (MHR (Fort Leonard Wood/Fort Jackson) = 1.7, 95 percent CI = 1.1–2.4); and slower 2-mile run times (MHR (Q4/Q1) = 1.8, 95 percent CI = 1.2–2.7).

(4) Traumatic Time-Loss Injury.

(a) For men, a higher risk of traumatic time-loss injury was associated with the Black and Native American races (MHR (Black/Caucasians) = 1.4, 95 percent CI = 1.0–2.0), MHR (Native American/Caucasian) = 1.8, 95 percent CI = 1.2–2.8); a current self-reported injury (MHR (yes/no) = 1.6, 95 percent CI = 1.1–2.4); and smoking before entering the Army (MHR (≤ 10 cigarettes/nonsmokers) = 1.5, 95 percent CI = 1.0–2.2, MHR (10–20 cigarettes/nonsmokers) = 1.4, 95 percent CI = 1.0–2.0, MHR (> 20 cigarettes/nonsmokers) = 1.7, 95 percent CI = 1.2–2.5).

(b) For women, a higher risk of traumatic time-loss injury was associated with a current self-reported injury (MHR (yes/no) = 2.2, 95 percent CI = 1.1–4.2).

4. DISCUSSION.

a. For men, injury risk was higher in smokers than nonsmokers in all four injury categories in consonance with previous studies in BCT. There was also a dose-response relationship showing that the risk of injury increases with the number of cigarettes smoked per day.

b. Injury risk for those with current self-reported injuries believed to adversely affect their AIT performance was approximately 2.2 times higher for men and 1.5 times higher for women (except in the traumatic injury category). Previous injury, both overuse and traumatic, put Soldiers at a higher risk for current injury. Civilian studies have also shown that those with a previous injury had a higher risk of reinjury than those who reported no previous injury.

5. CONCLUSIONS. This study identified risk factors for time-loss injury in Ordnance School AIT Soldiers. Overall, 31 percent of men and 54 percent of women involved in the project incurred at least one time-loss injury. When examining injury risk for all four injury categories, both cigarette use and self-reported injury were associated with a higher risk of injury in men. For three out of the four injury categories, self-reported injury (women), sit-ups (men) and the 2-mile run (men and women) were associated with a higher risk of injury.

6. RECOMMENDATIONS. In an effort to reduce injuries, surveillance and tracking of injuries in AIT Soldiers could alert commanders to elevated levels of injuries or to injury outbreaks. Smoking cessation classes and fitness training prior to entry are potential strategies to reduce injuries.

TABLE OF CONTENTS

1.	REFERENCES.....	1
2.	PURPOSE.....	1
3.	AUTHORITY.....	1
4.	BACKGROUND.....	1
5.	LITERATURE	2
	a. Injury Incidence and Injury Risk Factors in Basic Combat Training and Advanced Individual Training.....	2
	b. Overuse Injuries.....	3
	c. Traumatic Injuries	3
6.	METHODS	3
	a. Participants.....	3
	b. Questionnaires	4
	c. Army Physical Fitness Test Scores	4
	d. Injury Data	4
	e. Data Analysis	5
7.	RESULTS.....	6
	a. Descriptive Statistics	6
	b. Risk Factors for Time-Loss Injury	8
	c. Risk Factors for Overuse Time-Loss Injury.....	11
	d. Risk Factors for Lower Extremity Overuse Time-Loss Injury.....	14
	e. Risk Factors for Traumatic Time-Loss Injury	17
8.	DISCUSSION.....	19
	a. General Findings	19
	b. Age.....	21
	c. Race	22
	d. Basic Training Site	23
	e. Military Occupational Specialty.....	24
	f. Self-Reported Injury.....	24
	g. Cigarette Use	25
	h. Smokeless Tobacco Use.....	26
	i. Muscular Endurance.....	27
	j. Two-Mile Run Times.....	27
9.	CONCLUSIONS	28
10.	RECOMMENDATIONS	28
11.	TECHNICAL ASSISTANCE	28

Appendices

A.	REFERENCES	A-1
B.	SOLDIER HEALTH INPROCESSING QUESTIONNAIRE.....	B-1
C.	INJURY SHEET	C-1

List of Tables

1.	Cumulative Incidence of Injury and Injury Incidence Rates During Army Basic Training	2
2.	Military Occupational Specialties at Aberdeen Proving Ground	4
3.	Advanced Individual Training Ordnance School Soldier Health In-Processing Questionnaire Variables (Descriptive Statistics)	7
4.	Army Physical Fitness Test Scores	8
5.	Person-Time Injury Incidence Rates for All Four Injury Outcomes (Injuries/10,000 Person-Days)	8
6.	Univariate Cox Regression: Risk Factors Associated with Time- Loss Injuries in Ordnance Advanced Individual Training Subjects	9
7.	Multivariate Cox Regression: Risk Factors Associated with Time- Loss Injuries in Ordnance Advanced Individual Training Subjects	11
8.	Univariate Cox Regression: Risk Factors Associated with Overuse Time-Loss Injuries in Ordnance Advanced Individual Training Subjects.....	12
9.	Multivariate Cox Regression: Risk Factors Associated with Overuse Time-Loss Injuries in Ordnance Advanced Individual Training Subjects.....	14
10.	Univariate Cox Regression: Risk Factors Associated with Lower Extremity Overuse Time-Loss Injuries in Ordnance Advanced Individual Training Subjects.....	15
11.	Multivariate Cox Regression: Risk Factors Associated with Lower Extremity Overuse Time-Loss Injuries in Ordnance Advanced Individual Training Subjects.....	16
12.	Univariate Cox Regression: Risk Factors Associated with Traumatic Time-Loss Injuries in Ordnance Advanced Individual Training Subjects.....	17
13.	Multivariate Cox Regression: Risk Factors Associated with Traumatic Time-Loss Injuries in Ordnance Advanced Individual Training Subjects.....	19
14.	Risk Factors From All Four Injury Categories Placing Soldiers at a Higher Risk for a Time-Loss Injury (Multivariate Analysis)	20

INJURY PREVENTION REPORT NO. 12-HF-0893-10
INJURY INCIDENCE AND INJURY RISK FACTORS AMONG SOLDIERS IN THE
U.S. ARMY ORDNANCE SCHOOL
ABERDEEN PROVING GROUND, MARYLAND
JANUARY 2000–JUNE 2003

1. REFERENCES. Appendix A contains a listing of the references used in this report.
2. PURPOSE. From January 2000 to June 2003, the U.S. Army Public Health Command (Provisional) (USAPHC (Prov)), formerly known as the U.S. Army Center for Health Promotion and Preventive Medicine (USACHPPM), and the Kirk U.S. Army Health Clinic collaborated with the 143rd Ordnance Battalion at Aberdeen Proving Ground (APG), Maryland, to identify injuries and injury risk factors during advanced individual training (AIT). This report provides the results of this collaboration by addressing injury rates and injury risk factors among Soldiers in AIT at the Ordnance School.
3. AUTHORITY. Under Army Regulation (AR) 40-5 (paragraph 2-19), USACHPPM is responsible for providing support to Army preventive medicine activities, to include interpreting surveillance data, identifying leading health problems, and assisting in prevention and control of leading health problems.¹ This project was conducted under a cooperative agreement among the 61st Ordnance Brigade, Kirk U.S. Army Health Clinic, and USACHPPM, all located at APG.
4. INTRODUCTION. Time lost from work and training due to injuries can result in decreased military readiness and can compromise mission accomplishment. In 2004, Department of Defense (DOD) Service members experienced almost 25 million days of limited duty due to injuries. The top five injuries ranked by the number of days of limited duty were lower extremity overuse (pain, inflammation and stress fractures); lower extremity fractures; upper extremity fractures; torso overuse (pain, inflammation, and stress fractures); and lower extremity sprains and strains.² In a U.S. Marine Corps basic training study, investigators estimated that injuries among 22,000 male recruits resulted in more than 53,000 lost training days at a cost of \$16.5 million (1993 dollars).³ Other researchers examining infantry Soldiers found that fractures resulted in an average of 103 days of limited duty, sprains an average of 17 days of limited duty, tendinitis an average of 7 days of limited duty, and strains and musculoskeletal pain an average of 3 days of limited duty.⁴ In basic combat training (BCT), the incidence of injury has ranged from 21 percent to 42 percent for men and from 41 percent to 67 percent for women.⁵ There are only two previous reports on injuries and injury risk factors among Soldiers in AIT. One study involved medic AIT⁶ and the other investigation was an abstract reporting the preliminary results from the present project.⁷

5. BACKGROUND LITERATURE .

a. Injury Incidence and Injury Risk Factors in Basic and Advanced Individual Training.

(1) Cumulative injury incidence (proportion of trainees who experience one or more injuries during training) and injury rates (injured trainees per month) have been examined in the basic training units of the Army as well as in an Army AIT investigation.^{6,8–22} These data are shown in Table 1. In October 1998, Army BCT was extended from 8 to 9 weeks; therefore studies performed before and after this time are designated in Table 1 to reflect the increased time Soldiers were at risk in the investigations subsequent to October 1998.

Table 1. Cumulative Incidence of Injury and Injury Incidence Rates During Army Training

	Length of Training (weeks)	Study (Reference Number)	Year Data Collected	Recruits (n)		Cumulative Injury Incidence (%)		Injury Incidence Rate (%/month)	
				Men	Women	Men	Women	Men	Women
Army Basic Combat Training	8 weeks	12 ^a	1980	1,840	644	20.7	41.2	10.4	20.6
		11	1984	124	186	27.4	50.5	13.7	25.3
		18	1988	509	352	27.0	57.0	13.5	28.5
		16	1994	ND ^b	165	ND ^b	66.7	ND ^b	33.3
		19	1996	159	84	41.5	65.5	20.8	32.8
		13	1998	604	305	30.8	58.0	15.4	29.0
	9 weeks	20	1998	655	498	29.98	65.3	13.3	29.0
		6	2000	371	237	26.1	51.5	11.6	22.9
		21 ^c	2000	682/441	579/554	13.5/16.9	36.1/46.8	6.0/7.5	16.0/20.8
		22 ^{cd}	2003	442/569	295/377	19.5/27.9	41.0/47.7	8.7/12.4	18.2/21.2
		17	2007	2,147	915	36.9	64.7	16.4	28.8
Medic Advanced Individual Training	10 weeks	6	2000	439	287	24.0	30.0	9.6	12.0
Notes: ^a Injury data from self-report questionnaire ^b ND=No data collected on other gender ^c Cohort study with two groups ^d Injury data from surveillance system									

(2) In addition to cumulative injury incidence and injury rates, injury risk factors have been identified. Injury risk factors identified during basic training included female

gender, low aerobic fitness, cigarette smoking prior to BCT, low physical activity prior to basic training, low muscular endurance, and training in the summer compared to training in the fall.^{8–16, 23–27} A majority of the injuries occurring in Service members can be classified as either overuse or traumatic injury.²

b. Overuse Injuries. Overuse injuries are the result of abnormal and repetitive stress resulting in microtrauma to the soft tissues, bones or joints.²⁸ Some examples of overuse injuries include shin splints, tendonitis, stress fractures and bursitis.²⁹ In a study investigating injuries occurring in Army BCT and AIT, investigators found the largest proportion of injuries in both BCT and AIT were of the overuse type and involved the lower body.⁶ In a study of Marine Corps basic training, investigators found that approximately 80 percent of injuries were the overuse type and involved the lower extremities (primarily the knee and ankle/foot regions).³

c. Traumatic Injuries. Traumatic injuries are a result of an outside agent or force that causes acute damage or harm to the structure or function of the body.²⁸ Some examples of traumatic injuries include contusions, fractures, joint dislocations, concussions, strains and sprains.²⁹ In athletes the majority of traumatic injuries relates to overstretching of the soft tissue.³⁰ When soft tissue is suddenly stretched beyond its yield point, it will tear or rupture. The muscles most prone to overstretching or tearing are those that cross two joints.³⁰ Risk factors associated with traumatic injuries include previous injury, tobacco use, and strength imbalances. In regard to previous injuries, an investigation found that basketball players with a history of ankle injuries were 5 times more likely to sustain another ankle injury.³¹ For tobacco use, cigarette smokers have been shown to have higher scores on various measures of risk-taking behaviors possibly placing them at a higher risk of incurring a traumatic injury.³² Muscle strength and balance abnormalities have also been investigated and associated with acute muscle injuries.^{33,34}

6. METHODS.

a. Participants. Participants were Army personnel Service members attending AIT at APG (Edgewood Area) from January 2000 to June 2003. Soldiers attending AIT in the Edgewood Area of APG trained to qualify for one of five different military occupational specialties (MOSSs) (Table 2). There were three companies (Alpha, Bravo, and Charlie) in the single training battalion at the Edgewood Area of APG (143rd Ordnance Battalion).

Table 2. Military Occupational Specialties at Aberdeen Proving Ground

Military Occupational Specialty	Weeks of Training
Track Vehicle Repairer (63H)	16
Wheeled Vehicle Repairer (63W)	13
Self Propelled Field Artillery System Mechanic (63D)	10
Fuel and Electrical System Repairer (63G)	9
Track Vehicle Mechanic (63Y)	12

b. Questionnaires. Upon arrival at APG, each Service member was asked to complete the Soldier health in-processing (SHIP) questionnaire (appendix B) as a part of the in-processing procedures. The questionnaire asked the Service members about their demographic and lifestyle characteristics. Information requested included rank, race, gender, date of birth, BCT site, whether or not the student currently had an injury or illness perceived to affect their AIT performance, and history of tobacco use.

c. Army Physical Fitness Test Scores. Army Physical Fitness Test (APFT) data were obtained from the 143rd Ordnance Battalion Training and Operations (S-3) office. The APFT consisted of three events: a 2-minute maximal effort push-up event, a 2-minute maximal effort sit-up event, and a 2-mile run performed for time. For the push-up, the subject lowered his or her body in a generally straight line to a point where his or her upper arm was parallel to the ground, and then returned to the starting point with elbows fully extended. For the sit-up, the subject's knees were bent at a 90° angle, fingers were interlocked behind the head, and a second person held the subject's ankles while the subject kept his or her heels firmly on the ground. The subject raised his upper body to a vertical position so that the base of the neck was anterior to the base of the spine and then returned to the starting position. Scores were the number of push-ups and sit-ups that were successfully completed within the separate 2-minute time periods. The performance measure for the run was the time taken to complete the 2-mile distance. Time between events was no less than 10 minutes.

d. Injury Data. Injuries were obtained from an injury surveillance system in the medical clinic serving the AIT Soldiers. Every time a Soldier reported to the medical clinic for an injury, a medical provider would fill out an injury sheet (appendix C). The medical provider could record the type of injury that occurred and the number of profile or quarter days given to the Soldier by checking the appropriate boxes on the injury sheet. Injuries from the data sheet were then classified into four categories: time-loss injuries, overuse time-loss injuries, lower extremity overuse time-loss injuries, and traumatic time-loss injuries. Time-loss injuries included an injury of any type (overuse, traumatic, other, unknown) plus a profile of 1 or more days. Overuse time-loss injuries were identified as such by the medical provider in item 4 (Injury Category) on the injury

sheet and included a profile of one or more days. Lower extremity overuse time-loss injuries were identified as overuse injuries on the injury sheet in item 4 (Injury Category) and included a profile of 1 or more days, but were limited to the injuries occurring to the leg (upper and lower), knee, ankle and foot. Traumatic time-loss injuries were identified as such on the injury sheet in item 4 (Injury Category) and included a profile of 1 or more days.

e. Data Analysis.

(1) The questions about tobacco use on the SHIP survey (appendix B) asked if the Service member had smoked one or more cigarettes within the 30 days prior to BCT and if he or she had smoked on 20 of the 30 days prior to BCT. If Soldiers answered “yes” to smoking one or more cigarettes within the last 30 days prior to BCT, but “no” to the question asking if they had smoked on 20 or more days in the 30 days prior to BCT, they were considered an “occasional smoker.” If they answered yes to smoking on 20 of the last 30 days prior to BCT, they were considered a “frequent smoker.” Those who answered “yes” to using smokeless tobacco at least once in the 30 days prior to BCT, but “no” to the question asking if they had used smokeless tobacco on 20 or more days in the 30 days prior to BCT, were considered “occasional smokeless tobacco users,” and those who reported using smokeless tobacco on 20 or more days in the 30 days prior to BCT were considered “frequent smokeless tobacco users.”

(2) The age of the Soldier was determined by his or her response to question 9 on the SHIP survey (appendix B). Age was then grouped into three categories: age 17–19 years, 20–24 years, and 25+ years.

(3) Cumulative time-loss injury incidence rates were calculated as follows:

$$(\text{Soldiers with } \geq 1 \text{ time-loss injury}) / (\text{Total number of Soldiers in the Ordnance School}) \times 100\%$$

(4) Person-time injury incidence rates for each injury category were calculated as follows:

$$(\text{Soldiers with } \geq 1 \text{ time-loss injury}) / (\text{Total Soldier days in the Ordnance School} \times 10,000)$$

(5) The Statistical Package for the Social Sciences (SPSS), Version 16.0, was used for statistical analysis. Descriptive statistics (frequencies) were calculated for demographics (age, gender, race, military rank); BCT site; a current self-reported injury; a current self-reported illness; tobacco use variables; and push-ups, sit-ups, and the

2- mile run. Potential risk factors for time-loss injury, overuse time-loss injury, lower extremity overuse time-loss injury, and traumatic time-loss injury were explored using univariate and multivariate Cox regression. Hazard ratios and 95 percent confidence intervals (CIs) were calculated for each risk factor. Variables from the univariate analysis with a statistical significance of $p \leq 0.10$ were selected for a backward-stepping multivariate Cox regression. Since all of the tobacco variables were correlated (60 percent of smokeless users were also smokers), it was decided to only use the question, "How many cigarettes were smoked in the last 30 days before BCT," in the multivariate models. Multivariate hazard ratios (MHRs) and 95 percent CIs were calculated. For the Cox regression, APFT scores were converted to four quartiles (Qs) based on the distribution of scores where Q1 = high performance and Q4 = low performance.

7. RESULTS.

a. Descriptive Statistics.

(1) There were 3757 men and 498 women involved in the project. Table 3 displays the results from the SHIP questionnaire. A majority of the subjects were Caucasian men between the ages of 17 and 19 who had attended BCT at Fort Knox and were wheeled vehicle repairers (MOS 63W) of lower military rank (E1) and nonsmokers and non-smokeless tobacco users. The mean age (\pm standard deviation) for both men and women was 20 ± 3 years. Most of the women had attended BCT at Fort Jackson. A self-reported injury perceived to negatively affect AIT performance was reported by 7 percent of subjects, and a self-reported illness perceived to negatively affect AIT performance was reported by 2 percent of the subjects.

(2) When asked about smokeless tobacco use, men were 2.5 times more likely to be occasional smokeless tobacco users and 5.5 times more likely to be frequent smokeless tobacco users when compared to women. When asked if they had a current injury perceived to negatively affect AIT performance, women were 2.3 times more likely to answer yes when compared to men.

Injury Prevention Report No. 12-HF-0893-10, January 2000–June 2003

Table 3. Advanced Individual Training Ordnance School Soldier Health In-Processing Questionnaire Variables (Descriptive Statistics)

Variable	Level of Variables	Men n (%)	Women n (%)	Men and Women n (%)
Gender	Men Women Men and Women	3757 (100)	498 (100)	4255 (100)
Age	17–19 years 20–24 years 25+ years	2067 (55) 1324 (35) 366 (10)	242 (49) 171 (34) 85 (17)	2309 (54) 1495 (35) 451 (11)
Race	Caucasian Asian Black Hispanic Native	2452 (65) 103 (3) 480 (13) 506 (14) 216 (6)	330 (66) 9 (2) 87 (18) 44 (9) 28 (6)	2782 (65) 112 (3) 567 (13) 550 (13) 244 (6)
Rank	E1 E2 E3 E4+	2135 (57) 911 (24) 640 (17) 71 (2)	241 (48) 140 (28) 101 (20) 16 (3)	2376 (56) 1051 (25) 741 (17) 87 (2)
Basic Training Site	Fort Jackson Fort Knox Fort Leonard Wood Fort Benning Fort Sill	907 (24) 2135 (57) 189 (5) 357 (10) 169 (5)	401 (81) 0 (0) 69 (14) 0 (0) 28 (6)	1308 (31) 2135 (50) 258 (6) 357 (8) 197 (5)
Military Occupational Specialty	Track Vehicle Repairer (63H) Wheeled Vehicle Repairer (63W) Self-Propelled Field Artillery System Mechanic (63D) Fuel and Electrical System Repairer (63G) Track Vehicle Mechanic (63Y)	554 (15) 2168 (58) 214 (6) 286 (8) 535 (14)	59 (12) 352 (71) 0 (0) 28 (6) 59 (12)	613 (14) 2520 (59) 214 (5) 314 (7) 594 (14)
Self-Reported Injury	No Yes	3524 (94) 233 (6)	431 (87) 67 (14)	3955 (93) 300 (7)
Self-Reported Illness	No Yes Missing	3675 (98) 69 (2) 13 (0.3)	483 (97) 13 (3) 2 (0.4)	4158 (98) 82 (2) 15 (0.4)
Cigarette Use	Nonsmoker Occasional Frequent	2166 (58) 201 (5) 1390 (37)	296 (59) 34 (7) 168 (34)	2462 (58) 235 (6) 1558 (37)
How many cigarettes smoked per day?	Nonsmoker ≤ 10 10-20 ≥ 20 Missing	2166 (58) 367 (10) 582 (16) 408 (11) 234 (6)	296 (59) 65 (13) 58 (12) 40 (8) 39 (8)	2462 (58) 432 (10) 640 (15) 448 (11) 273 (6)
Smokeless Tobacco Use	Non-user Occasional smokeless Frequent smokeless	3158 (84) 171 (5) 428 (11)	481 (97) 8 (2) 9 (2)	3639 (86) 179 (4) 437 (10)
How much smokeless tobacco used per day?	Non-user ≤ 1 can 1 can or more ≥ 2 cans Missing	3158 (84) 262 (7) 122 (3) 27 (1) 188 (5)	481 (97) 6 (1) 2 (0.4) 0 (0) 9 (2)	3639 (86) 268 (6) 124 (3) 27 (1) 197 (5)

(3) Table 4 displays APFT scores for men and women. Compared to women, men performed an average of 22 more push-ups, an average of 2 more sit-ups, and ran the 2 miles 3.4 minutes faster.

Table 4. Army Physical Fitness Test Scores

Variable	Men n = 3757			Women n = 498		
	Mean \pm SD	Min	Max	Mean \pm SD	Min	Max
Push-Ups (Repetitions)	52 \pm 12	13	113	30 \pm 10	2	81
Sit-Ups (Repetitions)	62 \pm 10	22	109	60 \pm 12	26	122
2-Mile Run (Minutes)	14.9 \pm 1.5	10.9	32.9	18.3 \pm 2.0	13.6	30.6

(4) Table 5 displays person-time injury incidence rates for time-loss injury, overuse time-loss injury, lower extremity overuse time-loss injury, and traumatic time-loss injury for men and women. In all cases, injury rates are higher for men than women. Cumulative time-loss injury incidence was 31percent for men and 54 percent for women. When including both time-loss injuries and those not involving time loss, 36 percent of men and 61 percent of women experienced at least one injury during AIT.

Table 5. Person-Time Injury Incidence Rates for All Four Injury Outcomes (Injuries/10,000 Person-Days)

Injury Outcomes	Men Rate per 10,000 Person-Days	Women Rate per 10,000 Person-Days
Time-Loss Injuries	34.9	60.8
Overuse Time-Loss Injuries	23.8	46.8
Lower Extremity Overuse Time-Loss Injuries	20.7	44.1
Traumatic Time-Loss Injuries	8.9	11.7

b. Risk Factors for Time-Loss Injury.

(1) Table 6 displays the results of the univariate Cox regression with time-loss injury as the dependent variable. For men, time-loss injury risk was higher among those who were Native American (relative to Caucasian), of lower military rank (E1 relative to E2), who had attended BCT at Fort Jackson (relative to Fort Knox) and were smokers and/or smokeless tobacco users, with a self-reported injury, lower push-up

Injury Prevention Report No. 12-HF-0893-10, January 2000–June 2003

performance, lower sit-up performance, and slower two-mile run times. For women, time-loss injury risk was higher among those who were Caucasian (relative to Black), and frequent smokers and/or occasional smokeless tobacco users, with a self-reported injury, lower push-up performance, and slower 2-mile run times.

Table 6. Univariate Cox Regression: Risk Factors Associated with Time-Loss Injuries in Ordnance Advanced Individual Training

Variable	Men				Women			
	Variable Level	n (% TLI) ^a	Hazard Ratio (95% CI) ^b	p-value	Variable Level	n (% TLI) ^a	Hazard Ratio (95% CI) ^b	p-value
Age Group	17-19 years	2067 (30)	1.00		17-19 years	242 (54)	1.00	
	20-24 years	1324 (31)	1.05 (0.92-1.18)	0.48	20-24 years	171 (51)	0.89 (0.68-1.16)	0.39
	25+ years	366 (32)	1.10 (0.90-1.34)	0.37	25+ years	85 (61)	1.18 (0.86-1.63)	0.32
Race	Caucasian	2452 (32)	1.00		Caucasian	330 (57)	1.00	
	Asian	103 (25)	0.75 (0.51-1.1)	0.14	Asian	9 (44)	0.71 (0.26-1.91)	0.50
	Black	480 (33)	1.03 (0.87-1.22)	0.74	Black	87 (43)	0.69 (0.48-0.98)	0.04
	Hispanic	506 (24)	0.72 (0.59-0.87)	<0.01	Hispanic	44 (61)	1.18 (0.79-1.77)	0.42
	Native	216 (38)	1.26 (1.00-1.57)	0.05	Native	28 (54)	0.83 (0.49-1.41)	0.50
Rank	E1	2135 (32)	1.00		E1	241 (56)	1.00	
	E2	911 (28)	0.87 (0.75-1.00)	0.05	E2	140 (55)	1.02 (0.77-1.35)	0.88
	E3	640 (29)	0.92 (0.78-1.08)	0.31	E3	101 (48)	0.78 (0.56-1.09)	0.14
	E4+	71 (31)	0.92 (0.60-1.41)	0.69	E4+	16 (69)	1.36 (0.73-2.52)	0.33
Basic Training Site	Ft Jackson	907 (34)	1.00		Ft Jackson	401 (53)	1.00	
	Ft Knox	2135 (30)	0.86 (0.75-0.99)	0.03	Ft Knox	0		
	Ft Wood	189 (31)	0.90 (0.68-1.20)	0.48	Ft Wood	69 (59)	1.18 (0.85-1.65)	0.33
	Ft Benning	357 (31)	0.92 (0.74-1.14)	0.42	Ft Benning	0		
	Ft Sill	169 (27)	0.78 (0.57-1.06)	0.12	Ft Sill	28 (54)	1.03 (0.61-1.74)	0.91
Military Occupational Specialty	(63H)	554 (34)	1.00		(63H)	59 (63)	1.00	
	(63W)	2168 (30)	0.96 (0.81-1.13)	0.56	(63W)	352 (53)	0.98 (0.69-1.40)	0.91
	(63D)	214 (23)	0.97 (0.71-1.33)	0.85	(63D)	0	-	
	(63G)	286 (28)	1.14 (0.88-1.49)	0.33	(63G)	28 (32)	0.65 (0.31-1.36)	0.25
	(63Y)	535 (35)	1.13 (0.92-1.38)	0.25	(63Y)	59 (64)	1.25 (0.79-1.96)	0.34
Self-Reported Illness	No	3675 (31)	1.00		No	483 (54)	1.00	
	Yes	69 (33)	1.11 (0.73-1.67)	0.64	Yes	13 (62)	1.14 (0.57-2.31)	0.71
Self-Reported Injury	No	3524 (30)	1.00		No	431 (52)	1.00	
	Yes	233 (51)	2.27 (1.88-2.75)	<0.01	Yes	67 (66)	1.67 (1.21-2.30)	<0.01
Cigarette Use 30 Days Before BCT	Nonsmokers	2166 (26)	1.00		Nonsmokers	296 (52)	1.00	
	Occasional	201 (31)	1.19 (0.92-1.55)	0.19	Occasional	34 (56)	1.08 (0.67-1.73)	0.77
	Frequent	1390 (38)	1.56 (1.38-1.75)	<0.01	Frequent	168 (58)	1.27 (0.98-1.63)	0.07
How Many Cigarettes (cig) in 30 Days Before BCT?	(Nonsmokers)	2166 (26)	1.00		(Nonsmokers)	296 (52)	1.00	
	10 cig or less	367 (34)	1.29 (1.06-1.57)	0.01	10 cig or less	65 (52)	1.03 (0.71-1.49)	0.89
	10-20 cig	582 (37)	1.50 (1.28-1.75)	<0.01	10-20 cig	58 (62)	1.37 (0.95-1.97)	0.09
	20 cig or more	408 (45)	1.97 (1.67-2.33)	<0.01	20 cig or more	40 (65)	1.71 (1.12-2.59)	0.01
Smokeless Tobacco Use 30 Days Before BCT	Nonuser	3158 (30)	1.00		Nonuser	481 (54)	1.00	
	Occasional	171 (34)	1.13 (0.86-1.47)	0.38	Occasional	8 (75)	2.18 (0.97-4.90)	0.06
	Frequent	428 (37)	1.31 (1.11-1.55)	<0.01	Frequent	9 (67)	1.51 (0.67-3.38)	0.32
How Many Cans of Smokeless Tobacco 30 Days Before BCT?	0 (Nonuser)	3158 (30)	1.00		0 (Nonuser)	481 (54)	1.00	
	Less than 1	262 (37)	1.31 (1.06-1.61)	0.01	Less than 1	6 (50)	1.00 (0.32-3.14)	0.99
	1 on average	122 (39)	1.42 (1.06-1.89)	0.02	1 on average	2 (100)	2.17 (0.54-8.74)	0.28
	2 or more	27 (44)	1.59 (0.90-2.81)	0.11	2 or more	0 (0)		

Table 6. Univariate Cox Regression: Risk Factors Associated with Time-Loss Injuries in Ordnance Advanced Individual Training (continued)

Variable	Men				Women			
	Variable Level	n (% TLI) ^a	Hazard Ratio (95% CI) ^b	p-value	Variable Level	n (% TLI) ^a	Hazard Ratio (95% CI) ^b	p-value
Push-Ups (Repetitions)	0-43	963 (38)	1.63 (1.38-1.93)	<0.01	0-23	132 (60)	1.47 (1.03-2.09)	0.03
	44-50	967 (32)	1.38 (1.16-1.64)	<0.01	24-30	149 (58)	1.44 (1.02-2.04)	0.04
	50-59	915 (29)	1.22 (1.02-1.46)	0.03	31-36	106 (49)	1.04 (0.71-1.53)	0.84
	60+	912 (24)	1.00		37+	111 (47)	1.00	
Sit-Ups (Repetitions)	0-55	962 (34)	1.47 (1.24-1.75)	<0.01	0-53	131 (60)	1.28 (0.91-1.76)	0.16
	56-61	930 (34)	1.51 (1.27-1.80)	<0.01	54-60	125 (54)	1.17 (0.83-1.66)	0.36
	62-68	972 (31)	1.34 (1.12-1.59)	<0.01	61-67	120 (52)	1.04 (0.73-1.48)	0.82
	69+	893 (24)	1.00		68+	122 (50)	1.00	
2-Mile Run (Minutes)	0-13.91	921 (26)	1.00		0-17.00	126 (42)	1.00	
	13.92-14.77	979 (27)	1.02 (0.86-1.22)	0.80	17.01-18.08	124 (56)	1.46 (1.02-2.08)	0.04
	14.78-15.62	936 (31)	1.23 (1.03-1.46)	0.02	18.09-19.38	122 (51)	1.27 (0.88-1.83)	0.21
	15.63+	921 (40)	1.64 (1.39-1.93)	<0.01	19.39+	126 (68)	2.04 (1.45-2.88)	<0.01

^aTLI is time-loss injury. (Values are a percentage of the group that was injured.)^bCI is confidence interval.

(2) A backward-stepping multivariate analysis with time-loss injury as the dependent variable was performed with the following selected variables for men: race, rank, BCT site, self-reported injury, the number of cigarettes smoked in the 30 days before BCT, push-ups, sit-ups, and the 2-mile run. For men, rank, BCT, and push-ups did not reach the final step in the model. For women, the following variables were selected for inclusion in the multivariate model: race, self-reported injury, the number of cigarettes smoked in the 30 days before BCT, push-ups, and the 2-mile run. For women, the number of cigarettes smoked in the 30 days before BCT and push-ups did not reach the final step in the model.

(a) Table 7 displays the results of this analysis. For men, the risk of time-loss injury was independently associated with the Native American race (relative to Caucasians), self-reported injury, the number of cigarettes smoked in the 30 days before BCT, lower sit-up performance, and slower 2-mile run times. For women, a higher risk of time-loss injury was independently associated with the Caucasian race (relative to Blacks), self-reported injury, and slower 2-mile run times. Other multivariate models for men were also examined using the same variables above in the multivariate analysis.

(b) When a model was run limiting the fitness variables to just push-ups, push-ups made the final step and were found to place those who performed the least amount of push-ups at a higher risk of injury when compared to those who performed the most (MHR (Q4/Q1) = 1.4, 95 percent CI = 1.2-1.6). When another model was run substituting smokeless tobacco use for the number of cigarettes smoked per day, smokeless tobacco use made the final step, and frequent smokeless users were found

to have a higher risk of injury compared to non-smokeless users (MHR (frequent smokeless users/non-smokeless users) = 1.22, 95 percent CI = 1.03-1.45).

Table 7. Multivariate Cox Regression: Risk Factors Associated with Time-Loss Injuries in Ordnance Advanced Individual Training

Variable	Men (n=3523)				Women (n=459)			
	Variable Level	n	Hazard Ratio (95% CI) ^a	p-value	Variable Level	n	Hazard Ratio (95% CI) ^a	p-value
Race	Caucasian	2310	1.00		Caucasian	303	1.00	
	Asian	94	0.69 (0.45-1.06)	0.09	Asian	8	0.92 (0.34-2.48)	0.86
	Black	443	1.16 (0.96-1.39)	0.12	Black	81	0.65 (0.45-0.94)	0.02
	Hispanic	480	0.93 (0.76-1.14)	0.48	Hispanic	40	1.43 (0.93-2.19)	0.10
	Native	196	1.35 (1.06-1.71)	0.01	Native	27	0.77 (0.45-1.34)	0.36
Self-Reported Injury	No	3305	1.00		No	394	1.00	
	Yes	218	2.20 (1.80-2.67)	<0.01	Yes	65	1.60 (1.14-2.25)	<0.01
How Many Cigarettes 30 Days Before BCT	0 (Nonsmokers)	2166	1.00		b	b	b	b
	10 or less	367	1.25 (1.03-1.52)	0.03				
	10-20	582	1.46 (1.24-1.72)	<0.01				
	20 or more	408	1.87 (1.57-2.22)	<0.01				
Sit-Ups (Repetitions)	0-55	911	1.23 (1.02-1.48)	0.03	b	b	b	b
	56-61	865	1.35 (1.12-1.62)	<0.01				
	62-68	910	1.21 (1.01-1.46)	0.04				
	69+	837	1.00					
2-Mile Run (Minutes)	0-13.91	860	1.00		0-17.00	116	1.00	
	13.92-14.77	919	0.96 (0.80-1.15)	0.64	17.01-18.08	116	1.58 (1.09-2.30)	0.02
	14.78-15.62	876	1.07 (0.90-1.29)	0.44	18.09-19.38	112	1.32 (0.90-1.96)	0.16
	15.63+	868	1.41 (1.18-1.69)	<0.01	19.39+	115	2.17 (1.50-3.14)	<0.01

^aCI is confidence interval.

^bNot entered into the model because it did not meet the p<0.10 criteria in the univariate analysis.

c. Risk Factors for Overuse Time-Loss Injury.

(1) Table 8 displays the results of the univariate Cox regression with overuse time-loss injuries as the dependent variable. For men, overuse time-loss injury risk was higher among those who were Caucasian (relative to Hispanic); in the MOS of 63G (fuel and electrical system repairer) or 63Y (track vehicle mechanic) (relative to 63H (track vehicle repairer)); had a self-reported injury, smoked and/or used smokeless tobacco; and had a lower performance on push-ups, sit-ups, or the 2-mile run. For women, overuse time-loss injury risk was higher among those who were between 17 to 19 years old (relative to 20- to 24-year-olds), Caucasian (relative to Black), of lower military rank (E3 relative to E1), and who had attended basic training at Fort Leonard Wood (relative to Fort Jackson), had a self-reported injury, used smokeless tobacco frequently (relative to non-users), and had slower 2-mile run times.

Table 8. Univariate Cox Regression: Risk Factors Associated with Overuse Time-Loss Injuries in Ordnance Advanced Individual Training

Variable	Men				Women			
	Variable Level	n (% OI) ^a	Hazard Ratio (95% CI) ^b	p-value	Variable Level	n (% OI) ^a	Hazard Ratio (95% CI) ^b	p-value
Age Group	17-19 years	2067 (21)	1.00		17-19 years	242 (44)	1.00	
	20-24 years	1324 (21)	0.98 (0.84-1.14)	0.79	20-24 years	171 (36)	0.75 (0.55-1.03)	0.08
	25+ years	366 (20)	0.97 (0.75-1.24)	0.79	25+ years	85 (48)	1.11 (0.78-1.60)	0.56
Race	Caucasian	2452 (22)	1.00		Caucasian	330 (43)	1.00	
	Asian	103 (18)	0.76 (0.48-1.22)	0.26	Asian	9 (44)	1.01 (0.37-2.73)	0.99
	Black	480 (21)	0.92 (0.74-1.13)	0.42	Black	87 (32)	0.70 (0.47-1.05)	0.08
	Hispanic	506 (16)	0.69 (0.54-0.87)	<0.3	Hispanic	44 (48)	1.23 (0.78-1.95)	0.37
	Native	216 (25)	1.15 (0.86-1.52)		Native	28 (46)	1.03 (0.58-1.82)	0.92
Rank	E1	2135 (22)	1.00		E1	241 (43)	1.00	
	E2	911 (19)	0.87 (0.73-1.03)	0.11	E2	140 (44)	1.02 (0.74-1.40)	0.90
	E3	640 (20)	0.91 (0.75-1.10)	0.33	E3	101 (35)	0.71 (0.49-1.05)	0.08
	E4+	71 (20)	0.86 (0.50-1.46)	0.57	E4+	16 (56)	1.42 (0.72-2.81)	0.31
Basic Training Site	Ft Jackson	907 (21)	1.00		Ft Jackson	401 (40)	1.00	
	Ft Knox	2135 (21)	0.99 (0.84-1.18)	0.93	Ft Knox	0		
	Ft Wood	189 (18)	0.83 (0.58-1.20)	0.33	Ft Wood	69 (51)	1.40 (0.97-2.01)	0.08
	Ft Benning	357 (22)	1.05 (0.81-1.37)	0.71	Ft Benning	0		
	Ft Sill	169 (20)	0.93 (0.65-1.34)	0.71	Ft Sill	28 (43)	1.11 (0.62-2.00)	0.72
Military Occupational Specialty	(63H)	554 (22)	1.00		63H	59 (44)	1.00	
	(63W)	2168 (21)	1.10 (0.90-1.35)	0.35	63W	352 (42)	1.13 (0.74-1.72)	0.57
	(63D)	214 (16)	1.10 (0.75-1.61)	0.64	63D	0	-	
	(63G)	286 (19)	1.33 (0.96-1.83)	0.09	63G	28 (29)	0.86 (0.39-1.90)	0.70
	(63Y)	535 (25)	1.33 (1.04-1.70)	0.03	63Y	59 (46)	1.17 (0.68-2.01)	0.57
Self-Reported Illness	No	3675 (21)	1.00		No	483 (42)	1.00	
	Yes	69 (26)	1.26 (0.79-2.00)	0.34	Yes	13 (46)	1.19 (0.53-2.69)	0.67
Self-Reported Injury	No	3524 (20)	1.00		No	431 (40)	1.00	
	Yes	233 (37)	2.30 (1.84-2.87)	<0.01	Yes	67 (52)	1.56 (1.09-2.25)	0.02
Cigarette Use 30 Days Before BCT	Nonsmokers	2166 (17)	1.00		Nonsmokers	296 (40)	1.00	
	Occasional	201 (20)	1.16 (0.84-1.60)	0.38	Occasional	34 (38)	0.92 (0.52-1.63)	0.77
	Frequent	1390 (27)	1.67 (1.45-1.93)	<0.01	Frequent	168 (45)	1.22 (0.91-1.62)	0.18
How Many Cigarettes (cig) in 30 Days Before BCT?	Nonsmokers	2166 (17)	1.00		Nonsmokers	296 (40)	1.00	
	10 cig or less	367 (20)	1.17 (0.92-1.51)	0.21	10 cig or less	65 (43)	1.10 (0.73-1.65)	0.66
	10-20 cig	582 (28)	1.75 (1.46-2.10)	<0.01	10-20 cig	58 (48)	1.35 (0.89-2.04)	0.15
	20 cig or more	408 (32)	2.07 (1.70-2.52)	<0.01	20 cig or more	40 (50)	1.47 (0.92-2.37)	0.11
Smokeless Tobacco Use 30 Days Before BCT	Nonuser	3158 (20)	1.00		Nonuser	481 (41)	1.00	
	Occasional	171 (24)	1.17 (0.85-1.60)	0.35	Occasional	8 (63)	1.93 (0.79-4.68)	0.15
	Frequent	428 (28)	1.48 (1.21-1.79)	<0.01	Frequent	9 (67)	2.07 (0.92-4.68)	0.08
How Many Cans of Smokeless Tobacco 30 Days Before BCT?	Nonuser	3158 (20)	1.00		Nonuser	481 (41)	1.00	
	Less than 1	262 (29)	1.55 (1.22-1.97)	<0.01	Less than 1	6 (50)	1.40 (0.45-4.38)	0.56
	1 on average	122 (27)	1.41 (0.99-2.00)	0.06	1 on average	2 (100)	2.90 (0.72-11.68)	0.14
	2 or more	27 (33)	1.76 (0.91-3.40)	0.09	2 or more	0 (0)		
Push-Ups (Repetitions)	0-43	963 (27)	1.82 (1.48-2.23)	<0.01	0-23	132 (43)	1.23 (0.82-1.85)	0.31
	44-50	967 (23)	1.50 (1.22-1.85)	<0.01	24-30	149 (44)	1.30 (0.88-1.92)	0.20
	50-59	915 (18)	1.17 (0.93-1.46)	0.17	31-36	106 (43)	1.15 (0.75-1.76)	0.52
	60+	912 (16)	1.00		37+	111 (36)	1.00	

Table 8. Univariate Cox Regression: Risk Factors Associated with Overuse Time-Loss Injuries in Ordnance Advanced Individual Training (continued)

Variable	Men				Women			
	Variable Level	n (% OI) ^a	Hazard Ratio (95% CI) ^b	p-value	Variable Level	n (% OI) ^a	Hazard Ratio (95% CI) ^b	p-value
Sit-Ups (Repetitions)	0-55	962 (25)	1.67 (1.35-2.06)	<0.01	0-53	131 (47)	1.29 (0.88-1.90)	0.19
	56-61	930 (23)	1.58 (1.28-1.96)	<0.01	54-60	125 (44)	1.28 (0.89-1.90)	0.21
	62-68	972 (20)	1.35 (1.09-1.68)	<0.01	61-67	120 (38)	1.03 (0.69-1.56)	0.88
	69+	893 (16)	1.00		68+	122 (38)	1.00	
2-Mile Run (Minutes)	0-13.91	921 (17)	1.00		0-17.00	126 (33)	1.00	
	13.92-14.77	979 (19)	1.10 (0.88-1.36)	0.40	17.01-18.08	124 (39)	1.21 (0.80-1.83)	0.37
	14.78-15.62	936 (21)	1.29 (1.05-1.59)	0.02	18.09-19.38	122 (42)	1.30 (0.86-1.96)	0.21
	15.63+	921 (28)	1.73 (1.42-2.11)	<0.01	19.39+	126 (53)	1.87 (1.27-2.75)	<0.01

^aOI is overuse injury. (Values are a percentage of the group that was injured.)

^bCI is confidence interval.

(2) A backward-stepping multivariate analysis with overuse time-loss injury as the dependent variable was performed with the following selected variables for men: race, MOS, self-reported injury, the number of cigarettes smoked in the 30 days before BCT, push-ups, sit-ups, and the 2-mile run. For men, race and push-ups did not reach the final step in the model. For women, the following variables were selected for inclusion in the multivariate model: age, race, rank, basic training site, self-reported injury, and the 2-mile run. For women, rank did not reach the final step in the model.

(a) Table 9 displays the results of this analysis. For men, a higher risk of overuse time-loss injury was independently associated with MOS (63G (fuel and electrical system repairer) and 63Y (track vehicle mechanic) relative to MOS 63H (track vehicle repairer)); self-reported injury; the number of cigarettes smoked in the 30 days before BCT; lower sit-up performance; and slower 2-mile run times. For women, a higher risk of overuse time-loss injury was independently associated with 17- to 19-year-olds (relative to 20- to 24-year-olds), the Caucasian race (relative to Blacks), basic training attendance at Fort Leonard Wood (relative to Fort Jackson), a self-reported injury, and slower 2-mile run times. Other multivariate models for men were also examined using the same variables above in the multivariate analysis.

(b) When a model was run limiting the fitness variables to just push-ups, push-ups made the final step and were found to place those who performed the least amount of push-ups at a higher risk of injury when compared to those who performed the most (MHR (Q4/Q1) = 1.6, 95 percent CI = 1.3-1.9). When another model was run substituting smokeless tobacco use for the number of cigarettes smoked per day, smokeless tobacco use made the final step, and frequent smokeless users were found to have a higher risk of injury compared to non-smokeless users (MHR (frequent smokeless users/non-smokeless users) = 1.41, 95 percent CI = 1.16-1.71).

Table 9. Multivariate Cox Regression: Risk Factors Associated with Overuse Time-Loss Injuries in Ordnance Advanced Individual Training

Variable	Men (n=3523)				Women (n=498)			
	Variable Level	n	Hazard Ratio (95% CI) ^a	p-value	Variable Level	n	Hazard Ratio (95% CI) ^a	p-value
Age Group	b	b	b	b	17-19 20-24 25+	242 171 85	1.00 0.73 (0.53-1.00) 1.19 (0.81-1.74)	 0.05 0.39
Race	b	b	b	b	Caucasian Asian Black Hispanic Native	330 9 87 44 28	1.00 1.17 (0.43-3.19) 0.61 (0.40-0.92) 1.38 (0.86-2.20) 0.95 (0.53-1.70)	 0.76 0.02 0.18 0.86
Basic Training Site	b	b	b	b	Ft Jackson Ft Knox Ft Leonard Ft Benning Ft Sill	401 0 69 0 28	1.00 1.58 (1.08-2.31) 1.19 (0.66-2.15)	 0.02 0.57
Military Occupational Specialty	63H 63W 63D 63G 63Y	518 2033 204 264 504	1.00 1.17 (0.95-1.45) 1.13 (0.76-1.67) 1.50 (1.07-2.10) 1.35 (1.05-1.75)	 0.15 0.54 0.02 0.02	b	b	b	b
Self-Reported Injury	No Yes	3305 218	1.00 2.23 (1.76-2.81)	 <0.01	No Yes	431 67	1.00 1.47 (1.02-2.13)	 0.04
How Many Cigarettes 30 Days Before BCT	Nonsmokers 10 or less 10-20 20 or more	2166 367 582 408	1.00 1.12 (0.87-1.44) 1.68 (1.39-2.02) 1.93 (1.58-2.36)	 0.38 <0.01 <0.01	b	b	b	b
Sit-Ups (Repetitions)	0-55 56-61 62-68 69+	911 865 910 837	1.38 (1.10-1.73) 1.42 (1.13-1.78) 1.24 (0.99-1.56) 1.00	<0.01 <0.01 0.06	b	b	b	b
2-Mile Run (Minutes)	0-13.91 13.92-14.77 14.78-15.62 15.63+	860 919 876 868	1.00 1.00 (0.80-1.26) 1.14 (0.92-1.43) 1.46 (1.18-1.81)	 0.97 0.24 <0.01	0-17.00 17.01-18.08 18.09-19.38 19.39+	126 124 122 126	1.00 1.22 (0.81-1.86) 1.41 (0.93-2.13) 1.95 (1.31-2.90)	 0.34 0.11 <0.01

^aCI is confidence interval.
^bNot entered into the model because it did not meet the p<0.10 criteria in the univariate analysis

^aCI is confidence interval.^bNot entered into the model because it did not meet the p<0.10 criteria in the univariate analysis.d. Risk Factors for Lower Extremity Overuse Time-Loss Injury.

(1) Table 10 displays the results of the univariate Cox regression with lower extremity overuse time-loss injuries as the dependent variable. For men, lower extremity overuse time-loss injury risk was higher for those who were Caucasian (relative to Hispanic), in the MOS of 63Y (track vehicle mechanic) (relative to 63H (track vehicle repairer)), had a self-reported injury, smoked and/or used smokeless tobacco, and had a lower performance on push-ups, sit-ups, and slower 2-mile run times. For women, lower extremity overuse time-loss injury risk was higher for those who were of lower rank (E1 relative to E3), who had attended basic training at Fort Leonard Wood

Injury Prevention Report No. 12-HF-0893-10, January 2000–June 2003

(relative to attendance at Fort Jackson), had a self-reported injury, and had lower performance on push-ups and slower 2-mile run times.

Table 10. Univariate Cox Regression: Risk Factors Associated with Lower Extremity Overuse Time-Loss Injuries in Ordnance Advanced Individual Training

Variable	Men				Women			
	Variable Level	n (% LOI) ^a	Hazard Ratio (95% CI) ^b	p-value	Variable Level	n (% LOI) ^a	Hazard Ratio (95% CI) ^b	p-value
Age Group	17-19	2067 (18)	1.00		17-19	242 (41)	1.00	
	20-24	1324 (18)	1.01 (0.86-1.19)	0.91	20-24	171 (34)	0.80 (0.58-1.11)	0.18
	25+	366 (19)	1.06 (0.82-1.38)	0.64	25+	85 (47)	1.21 (0.83-1.74)	0.32
Race	Caucasian	2452 (19)	1.00		Caucasian	330 (40)	1.00	
	Asian	103 (17)	0.84 (0.52-1.36)	0.47	Asian	9 (44)	1.11 (0.41-3.00)	0.84
	Black	480 (17)	0.85 (0.67-1.07)	0.17	Black	87 (32)	0.77 (0.51-1.16)	0.21
	Hispanic	506 (14)	0.69 (0.54-0.89)	<0.01	Hispanic	44 (48)	1.36 (0.86-2.15)	0.19
	Native	216 (23)	1.22 (0.91-1.64)	0.18	Native	28 (39)	0.95 (0.51-1.76)	0.87
Rank	E1	2135 (19)	1.00		E1	241 (42)	1.00	
	E2	911 (17)	0.86 (0.72-1.04)	0.12	E2	140 (41)	0.94 (0.68-1.31)	0.73
	E3	640 (16)	0.86 (0.69-1.06)	0.15	E3	101 (30)	0.63 (0.42-0.95)	0.03
	E4+	71 (18)	0.92 (0.53-1.60)	0.76	E4+	16 (56)	1.46 (0.74-2.88)	0.28
Basic Training Site	Ft Jackson	907 (19)	1.00		Ft Jackson	401 (37)	1.00	
	Ft Knox	2135 (19)	0.99 (0.82-1.18)	0.88	Ft Knox	0 (0)		
	Ft Wood	189 (14)	0.73 (0.48-1.10)	0.13	Ft Wood	69 (51)	1.56 (1.08-2.25)	0.02
	Ft Benning	357 (18)	0.99 (0.75-1.32)	0.96	Ft Benning	0 (0)		
	Ft Sill	169 (18)	0.98 (0.67-1.43)	0.91	Ft Sill	28 (43)	1.22 (0.68-2.20)	0.50
Military Occupational Specialty	63H	554 (19)	1.00		63H	59 (44)	1.00	
	63W	2168 (18)	1.10 (0.88-1.37)	0.41	63W	352 (39)	1.02 (0.67-1.55)	0.94
	63D	214 (15)	1.16 (0.78-1.73)	0.47	63D	0 (0)		
	63G	286 (14)	1.07 (0.74-1.54)	0.71	63G	28 (29)	0.82 (0.37-1.81)	0.62
	63Y	535 (21)	1.26 (0.97-1.65)	0.09	63Y	59 (42)	1.04 (0.60-1.81)	0.89
Self-Reported Illness	No	3675 (18)	1.00		No	483 (39)	1.00	
	Yes	69 (22)	1.18 (0.71-1.97)	0.53	Yes	13 (46)	1.28 (0.57-2.88)	0.56
Self-Reported Injury	No	3524 (18)	1.00		No	431 (38)	1.00	
	Yes	233 (30)	2.06 (1.61-2.63)	<0.01	Yes	67 (49)	1.54 (1.06-2.24)	0.02
Cigarette Use 30 Days Before BCT	Nonsmokers	2166 (15)	1.00		Nonsmokers	296 (39)	1.00	
	Occasional	201 (18)	1.25 (0.89-1.76)	0.20	Occasional	34 (32)	0.83 (0.45-1.54)	0.55
	Frequent	1390 (24)	1.75 (1.50-2.04)	<0.01	Frequent	168 (42)	1.20 (0.89-1.62)	0.23
How Many Cigarettes (cig) in 30 Days Before BCT?	Nonsmokers	2166 (15)	1.00		Nonsmokers	296 (39)	1.00	
	10 cig or less	367 (17)	1.22 (0.93-1.60)	0.15	10 cig or less	65 (42)	1.13 (0.75-1.73)	0.56
	10-20 cig	582 (25)	1.78 (1.46-2.17)	<0.01	10-20 cig	58 (43)	1.26 (0.82-1.95)	0.30
	20 cig or more	408 (30)	2.24 (1.82-2.77)	<0.01	20 cig or more	40 (48)	1.46 (0.89-2.37)	0.13
Smokeless Tobacco Use 30 Days Before BCT	Non-user	3158 (17)	1.00		Non-user	481 (39)	1.00	
	Occasional	171 (22)	1.24 (0.88-1.73)	0.22	Occasional	8 (63)	2.03 (0.83-4.93)	0.12
	Frequent	428 (26)	1.57 (1.28-1.93)	<0.01	Frequent	9 (44)	1.29 (0.48-3.49)	0.61
How Many Cans of Smokeless Tobacco 30 Days Before BCT?	Non-use)	3158 (17)	1.00		Non-user	481 (39)	1.00	
	Less than 1	262 (26)	1.61 (1.25-2.07)	<0.01	Less than 1	6 (33)	0.89 (0.22-3.57)	0.87
	1 on average	122 (25)	1.55 (1.08-2.23)	0.02	1 on average	2 (50)	1.28 (0.18-9.11)	0.81
	2 or more	27 (30)	1.84 (0.91-3.69)	0.09	2 or more	0		
Push-Ups (Repetitions)	0-43	963 (23)	1.70 (1.36-2.12)	<0.01	0-23	132 (42)	1.38 (0.90-2.13)	0.14
	44-50	967 (20)	1.49 (1.19-1.86)	<0.01	24-30	149 (43)	1.50 (0.99-2.27)	0.06
	50-59	915 (16)	1.20 (0.94-1.52)	0.14	31-36	106 (41)	1.32 (0.84-2.07)	0.22
	60+	912 (14)	1.00		37+	111 (31)	1.00	

Table 10. Univariate Cox Regression: Risk Factors Associated with Lower Extremity Overuse Time-Loss Injuries in Ordnance Advanced Individual Training (continued)

Variable	Men				Women			
	Variable Level	n (% LOI) ^a	Hazard Ratio (95% CI) ^b	p-value	Variable Level	n (% LOI) ^a	Hazard Ratio (95% CI) ^b	p-value
Sit-Ups (Repetitions)	0-55	962 (22)	1.78 (1.41-2.23)	<0.01	0-53	131 (44)	1.37 (0.91-2.04)	0.13
	56-61	930 (20)	1.63 (1.29-2.06)	<0.01	54-60	125 (43)	1.39 (0.93-2.09)	0.11
	62-68	972 (18)	1.47 (1.16-1.86)	<0.01	61-67	120 (36)	1.09 (0.71-1.67)	0.70
	69+	893 (13)	1.00		68+	122 (34)	1.00	
2-Mile Run (Minutes)	0-13.91	921 (14)	1.00		0-17.00	126 (32)	1.00	
	13.92-14.77	979 (16)	1.11 (0.87-1.40)	0.40	17.01-18.08	124 (38)	1.23 (0.80-1.87)	0.35
	14.78-15.62	936 (18)	1.28 (1.02-1.61)	0.03	18.09-19.38	122 (38)	1.23 (0.80-1.87)	0.35
	15.63+	921 (25)	1.83 (1.47-2.26)	<0.01	19.39+	126 (50)	1.77 (1.19-2.64)	<0.01

^aLOI is lower extremity overuse injury. (Values are a percentage of the group that was injured.)
^bCI is confidence interval.

(2) A backward-stepping multivariate analysis with lower extremity overuse time-loss injury as the dependent variable was performed with the following selected variables for men: race, MOS, self-reported injury, the number of cigarettes smoked in the 30 days before BCT, push-ups, sit-ups, and 2-mile run times. For men, race, MOS, and push-ups did not reach the final step in the model. For women, the following variables were selected for inclusion in the multivariate model: rank, basic training site, self-reported injury, push-ups, and 2-mile run times. For women, rank and push-ups did not reach the final step in the model. Table 11 displays the results of this analysis. For men, a higher risk of lower extremity overuse time-loss injury was independently associated with self-reported injury, the number of cigarettes smoked in the 30 days before BCT, lower sit-up performance, and slower 2-mile run times. For women, a higher risk of lower extremity overuse time-loss injury was independently associated with those who had attended basic training at Fort Leonard Wood (relative to Fort Jackson), had a self-reported injury, and slower 2-mile run times.

Table 11. Multivariate Cox Regression: Risk Factors Associated with Lower Extremity Overuse Time-Loss Injuries in Ordnance Advanced Individual Training

Variable	Men (n=3523)				Women (n=498)			
	Category of Variable	n	Hazard Ratio (95% CI) ^a	p-value	Category of Variable	n	Hazard Ratio (95% CI) ^a	p-value
Basic Training Site					Ft Jackson	401	1.00	
					Ft Knox	0		
					Ft Leonard	69	1.67 (1.14-2.43)	<0.01
					Ft Benning	0		
Self-Reported Injury	No	3305	1.00		No	431	1.00	
	Yes	218	1.95 (1.51-2.52)	<0.01	Yes	67	1.39 (0.95-2.03)	0.09
How Many Cigarettes 30 Days Before BCT	Nonsmokers	2166	1.00					
	10 or less	367	1.17 (0.89-1.53)	0.25				
	10-20	582	1.69 (1.38-2.06)	<0.01				
	20 or more	408	2.08 (1.68-2.57)	<0.01				

Table 11. Multivariate Cox Regression: Risk Factors Associated with Lower Extremity Overuse Time-Loss Injuries in Ordnance Advanced Individual Training (continued)

Variable	Men (n=3523)				Women (n=498)			
	Category of Variable	n	Hazard Ratio (95% CI) ^a	p-value	Category of Variable	n	Hazard Ratio (95% CI) ^a	p-value
Sit-Ups (Repetitions)	0-55	911	1.45 (1.13-1.85)	<0.01	b	b	b	b
	56-61	865	1.42 (1.11-1.82)	<0.01				
	62-68	910	1.33 (1.04-1.70)	0.02				
	69+	837	1.00					
2-Mile Run (Minutes)	0-13.91	860	1.00		0-17.00	126	1.00	
	13.92-14.77	919	0.98 (0.77-1.25)	0.87	17.01-18.08	124	1.18 (0.77-1.80)	0.45
	14.78-15.62	876	1.09 (0.86-1.39)	0.48	18.09-19.38	122	1.23 (0.81-1.89)	0.34
	15.63+	868	1.49 (1.18-1.87)	<0.01	19.39+	126	1.80 (1.20-2.69)	<0.01

^aCI is confidence interval.
^bNot entered into the model because it did not meet the p<0.10 criteria in the univariate analysis.

e. Risk Factors for Traumatic Time-Loss Injury.

(1) Table 12 displays the results of the univariate Cox regression with traumatic time-loss injury as the dependent variable. For men, traumatic time-loss injury was higher among those who were Native American (relative to Caucasian), attended BCT at Ft Leonard Wood (compared to Ft Jackson), had a self-reported injury, smoked, used ≥ 2 cans of smokeless tobacco in the 30 days prior to BCT, and had a lower performance on sit-ups. For women, traumatic time-loss injury risk was higher among those who were Black (relative to Caucasian), had a self-reported injury, and smoked.

Table 12. Univariate Cox Regression: Risk Factors Associated with Traumatic Time-Loss Injuries in Ordnance Advanced Individual Training

Variable	Men				Women			
	Category of Variable	n (% TI) ^a	Hazard Ratio (95% CI) ^b	p-value	Category of Variable	n (% TI) ^a	Hazard Ratio (95% CI) ^b	p-value
Age Group	17-19	2067 (8)	1.00		17-19	242 (10)	1.00	
	20-24	1324 (8)	0.97 (0.76-1.24)	0.82	20-24	171 (12)	1.39 (0.76-2.52)	0.29
	25+	366 (7)	0.81 (0.53-1.24)	0.34	25+	85 (9)	1.03 (0.46-2.31)	0.95
Race	Caucasian	2452 (8)	1.00		Caucasian	330 (13)	1.00	
	Asian	103 (7)	0.90 (0.42-1.91)	0.78	Asian	9 (0)	c	
	Black	480 (9)	1.26 (0.91-1.74)	0.17	Black	87 (5)	0.35 (0.12-0.97)	0.04
	Hispanic	506 (6)	0.80 (0.54-1.16)	0.24	Hispanic	44 (5)	0.35 (0.08-1.43)	0.14
	Native	216 (12)	1.55 (1.02-2.36)	0.04	Native	28 (7)	0.50 (0.12-2.07)	0.34
Rank	E1	2135 (8)	1.00		E1	241 (12)	1.00	
	E2	911 (9)	1.12 (0.86-1.46)	0.42	E2	140 (10)	0.83 (0.47-1.68)	0.71
	E3	640 (7)	0.97 (0.70-1.34)	0.84	E3	101 (8)	0.68 (0.31-1.49)	0.34
	E4+	71 (4)	0.45 (0.14-1.44)	0.18	E4+	16 (6)	0.56 (0.08-4.15)	0.57
Basic Training Site	Ft Jackson	907 (7)	1.00		Ft Jackson	401 (11)	1.00	
	Ft Knox	2135 (8)	1.14 (0.86-1.52)	0.37	Ft Knox	0 (0)	--	
	Ft Leonard	189 (11)	1.57 (0.95-2.59)	0.08	Ft Leonard	69 (7)	0.65 (0.26-1.64)	0.36
	Ft Benning	357 (9)	1.28 (0.83-1.97)	0.26	Ft Benning	0 (0)	--	
	Ft Sill	169 (5)	0.79 (0.39-1.58)	0.50	Ft Sill	28 (18)	1.93 (0.76-4.90)	0.17

Table 12. Univariate Cox Regression: Risk Factors Associated with Traumatic Time-Loss Injuries in Ordnance Advanced Individual Training (continued)

Variable	Men				Women			
	Category of Variable	n (% TI) ^a	Hazard Ratio (95% CI) ^b	p-value	Category of Variable	n (% TI) ^a	Hazard Ratio (95% CI) ^b	p-value
Military Occupational Specialty	63H	554 (9)	1.00		63H	59 (15)	1.00	
	63W	2168 (8)	1.10 (0.79-1.52)	0.58	63W	352 (11)	0.96 (0.45-2.08)	0.92
	63D	214 (7)	1.30 (0.72-2.34)	0.38	63D	0 (0)	--	
	63G	286 (4)	0.73 (0.38-1.38)	0.33	63G	28 (4)	0.49 (0.06-3.89)	0.48
	63Y	535 (8)	0.97 (0.63-1.47)	0.87	63Y	59 (9)	0.74 (0.24-2.25)	0.74
Self-Reported Illness	No	3675 (8)	1.00		No	483 (10)	1.00	
	Yes	69 (7)	0.89 (0.37-2.15)	0.79	Yes	13 (15)	1.58 (0.38-6.49)	0.53
Self-Reported Injury	No	3524 (8)	1.00		No	431 (9)	1.00	
	Yes	233 (12)	1.59 (1.07-2.36)	0.02	Yes	67 (18)	2.19 (1.14-4.20)	0.02
Cigarette Use 30 Days Before BCT	Nonsmokers	2166 (7)	1.00		Nonsmokers	296 (10)	1.00	
	Occasional	201 (9)	1.32 (0.81-2.15)	0.27	Occasional	34 (6)	0.61 (0.15-2.59)	0.51
	Frequent	1390 (10)	1.44 (1.14-1.83)	<0.01	Frequent	168 (13)	1.38 (0.79-2.43)	0.26
How Many Cigarettes (cig) in 30 Days Before BCT?	Nonsmokers	2166 (7)	1.00		Nonsmokers	296 (10)	1.00	
	10 cig or less	367 (10)	1.48 (1.03-2.13)	0.04	10 or less	65 (3)	0.32 (0.08-1.35)	0.12
	10-20 cig	582 (9)	1.33 (0.97-1.83)	0.08	10-20	58 (17)	1.90 (0.92-3.91)	0.08
	20 cig or more	408 (11)	1.65 (1.18-2.31)	<0.01	20 or more	40 (20)	2.42 (1.10-5.33)	0.03
Smokeless Tobacco Use 30 Days Before BCT	Nonuser	3158 (8)	1.00		Nonuser	481 (10)	1.00	
	Occasional	171 (9)	1.05 (0.62-1.78)	0.86	Occasional	8 (25)	3.02 (0.73-12.45)	0.19
	Frequent	428 (9)	1.21 (0.87-1.70)	0.26	Frequent	9 (0)	c	
How Many Cans of Smokeless Tobacco 30 Days Before BCT?	Nonuser	3158 (8)	1.00		Nonuser	481 (10)		
	Less than 1	262 (8)	1.06 (0.68-1.66)	0.79	Less than 1	6 (0)	c	
	1 on average	122 (10)	1.30 (0.73-2.33)	0.37	1 on average	2 (0)	c	
	2 or more	27 (22)	3.24 (1.44-7.29)	<0.01	2 or more	0		
Push-Ups (Repetitions)	0-43	963 (8)	1.04 (0.76-1.43)	0.81	0-23	132 (14)	1.54 (0.72-3.31)	0.27
	44-50	967 (8)	0.98 (0.71-1.36)	0.93	24-30	149 (11)	1.22 (0.56-2.66)	0.62
	50-59	915 (7)	0.92 (0.66-1.29)	0.63	31-36	106 (5)	0.49 (0.17-1.44)	0.20
	60+	912 (8)	1.00		37+	111 (10)	1.00	
Sit-Ups (Repetitions)	0-55	962 (9)	1.29 (0.92-1.81)	0.14	0-53	131 (14)	1.76 (0.79-3.96)	0.17
	56-61	930 (9)	1.40 (1.00-1.96)	0.05	54-60	125 (11)	1.54 (0.67-3.56)	0.31
	62-68	972 (7)	1.13 (0.80-1.60)	0.50	61-67	120 (9)	1.28 (0.53-3.10)	0.58
	69+	893 (7)	1.00		68+	122 (7)	1.00	
2-Mile Run (Minutes)	0-13.91	921 (7)	1.00		0-17.00	126 (7)	1.00	
	13.92-14.77	979 (6)	0.87 (0.62-1.23)	0.43	17.01-18.08	124 (11)	1.53 (0.66-3.54)	0.32
	14.78-15.62	936 (8)	1.12 (0.80-1.55)	0.51	18.09-19.38	122 (10)	1.31 (0.55-3.12)	0.54
	15.63+	921 (10)	1.30 (0.93-1.76)	0.13	19.39+	126 (14)	1.70 (0.75-3.86)	0.21

^aTI is traumatic injury. (Values are a percentage of the group that was injured.)
^bCI is confidence interval.
^cWomen had no reported traumatic injuries.

(2) A backward-stepping multivariate analysis with traumatic time-loss injury as the dependent variable was performed with the following selected variables for men: race, basic training site, self-reported injury, the number of cigarettes smoked in the 30 days before BCT, and sit-ups. For men, the BCT site and sit-ups did not reach the final step in the model. For women, the following variables were selected for inclusion in the multivariate model: race and self-reported injury. For women, race did not reach the

final step in the model. Table 13 displays the results of this analysis. For men, a higher risk of traumatic time-loss injury was independently associated with Black and Native American race (relative to Caucasian), a self-reported injury, and smoking. For women, a higher risk of traumatic time-loss injury was independently associated with a self-reported injury.

Table 13. Multivariate Cox Regression: Risk Factors Associated with Traumatic Time-Loss Injuries in Ordnance Advanced Individual Training

Variable	Men (n=3523)				Women (n=498)			
	Category of Variable	n	Hazard Ratio (95% CI) ^a	p-value	Category of Variable	n	Hazard Ratio (95% CI) ^a	p-value
Race	Caucasian	2310	1.00		b	b	b	b
	Asian	94	0.91 (0.40-2.06)	0.82				
	Black	443	1.42 (1.00-2.03)	0.05				
	Hispanic	480	0.96 (0.65-1.43)	0.85				
	Native	196	1.81 (1.19-2.76)	<0.01				
Self-Reported Injury	No	3305	1.00		No Yes	431 67	1.00 2.19 (1.14-4.20)	0.02
	Yes	218	1.59 (1.06-2.38)	0.03				
How Many Cigarettes 30 Days Before BCT	Nonsmokers	2166	1.00		c	c	c	c
	10 or less	367	1.49 (1.03-2.15)	0.03				
	10-20	582	1.41 (1.02-1.96)	0.04				
	20 or more	408	1.74 (1.23-2.47)	<0.01				
^a CI is confidence interval.								
^b Did not reach the final step in the backward-stepping multivariate analysis.								
^c Not entered into the model because it did not meet the p<0.10 criteria in the univariate analysis.								

8. DISCUSSION.

a. General Findings.

(1) One of the main findings of the present study was that self-reported injury, cigarette use, and the 2-mile run were associated with a higher risk of injury in all four of the injury categories (except for the 2-mile run in the traumatic injuries category). For women, self-reported injury and slower 2-mile run times were associated with a higher risk of injury in three of the four injury categories. Table 14 displays the variables from all four injury categories (multivariate Cox regression analysis) associated with a higher risk of injury for men and women. It is interesting that only men were associated with a higher risk of injury relative to cigarette use and lower sit-up performance, and only women were associated with a higher risk of injury relative to attendance at basic training at Fort Leonard Wood. In most categories (excluding traumatic time loss), both men and women had higher injury risk if they had a self-reported injury or lower performance on the 2-mile run.

Injury Prevention Report No. 12-HF-0893-10, January 2000–June 2003

Table 14. Risk Factors From All Four Injury Categories Placing Soldiers at a Higher Risk for a Time-Loss Injury (Multivariate Analysis)

Variables	Variable Level	Men				Women			
		Time Loss	Overuse Time Loss	Lower Extremity Overuse Time Loss	Traumatic Time Loss	Time Loss	Overuse Time Loss	Lower Extremity Overuse Time Loss	Traumatic Time Loss
Age Group							X		
	17-19						X		
	20-24								
	25+								
Military Occupational Specialty			X						
	Track Vehicle Repairer (63H)								
	Wheeled Vehicle Repairer (63W)								
	Self-Propelled Field Artillery System Mechanic (63D)								
	Fuel and Electrical System Repairer (63G)		X						
	Track Vehicle Mechanic (63Y)		X						
Race		X			X	X			
	Caucasian								
	Asian								
	Black				X	X			
	Hispanic								
	Native	X			X				
Basic Training Site									
	Ft Jackson								
	Ft Knox								
	Ft Wood							X	
	Ft Benning								
	Ft Sill								
Self-Reported Injury		X	X	X	X	X	X		X
Cigarette Use		X	X	X	X				
	Nonsmoker								
	>10	X			X				
	10-20	X	X	X	X				
	20 or more	X	X	X	X				
Sit-Ups		X	X						
2-Mile Run		X	X	X		X	X	X	

(2) The relative risk for time-loss injuries was 1.74 times higher for women when compared to men, which is in agreement with studies performed during BCT.^{9,11} For time-loss injuries, the findings were injury incidence rates of 34.9/10,000 person-days for men and 60.8/10,000 person-days for women. A comparison of the rates found in

an investigation of the Army Combat Medic AIT course showed that men had a similar injury incidence rate of 34.2/10,000 person-days which was about the same as the current study. However, women in the Army Combat Medic course had an injury incidence rate of 42.8/10,000 person-days which is much lower than the Ordnance School injury incidence rate of 60.8/10,000 person-days.⁶ The rates in the Army Combat Medic AIT course were calculated assuming everyone completed the 10-week course. The higher injury incidence rates seen for women in the Ordnance School could be a result of the different types of training involved in the Ordnance School curriculum compared to the Army Combat Medic AIT course. In the Ordnance School, women are required to perform mechanical work on heavy equipment which requires more muscle mass and strength when compared to the training involved in the Army Combat Medic course. Also, when examining tobacco use for women, 34 percent of those in the Army Combat Medic AIT reported tobacco use within the last year compared to 41 percent of women in the current study who reported using tobacco (cigarettes) within the last 30 days. The specificity of the training in the Ordnance School requiring greater amounts of strength and the increase in tobacco use could be partially responsible for the higher injury incidence in women in the Ordnance School when compared to Army Combat Medic AIT.

b. Age.

(1) Younger age (17-19 years) was independently associated with a higher risk of injury for women when compared to slightly older women (20-24 years). Other investigations during BCT and AIT have shown that older recruits are at a higher risk of being injured.^{6,10,13,35} It has been suggested that when younger and older trainees all train at similar frequencies, intensities, and durations (as in AIT), the older trainees are at a greater risk of injury because of age-related fitness factors.¹⁰ With aging, there is a decrease in run speed and muscular endurance (which occurs around 30 years old), in addition to a decrease in lung vital capacity and aerobic capacity. These declines may contribute to the higher likelihood of injury.^{4,36}

(2) The civilian literature is inconsistent when investigating the association between age and injury, with some studies of physically active individuals showing no association, while other studies indicated that older age was associated with injury.³⁷⁻⁴³ Therefore it is unclear why the younger age group would be at a higher risk of injury when compared to a slightly older group who would also not be affected by any age-related fitness factors. Although when examining self-reported injury that may have an effect on AIT performance, 48 percent of 17-19-year-olds answered yes to this question compared to 33 percent of 20-24-year-olds.

c. Race.

(1) Native American men had a higher injury risk for time-loss injuries and traumatic time-loss injuries compared to Caucasians. One explanation could be that Native Americans have been shown to have a higher prevalence of tobacco use (cigarettes) compared to other ethnic groups.^{44,45} Tobacco use (cigarette smoking) has been shown to increase injury rates in BCT.^{13,35,46-48} However, in the current study the prevalence of cigarette use was higher in Caucasians (50 percent) compared to Native Americans (41 percent).

(a) When examining the reporting of no leisure time physical activity and ethnicity, one study found that 46 percent of Native Americans engaged in no leisure time physical activity compared to 36 percent of Caucasians, while another study found 33 percent of Native Americans report no leisure time physical activity compared to 28 percent of other racial/ethnic groups.^{45,49} However, physical activity levels in an AIT group would be similar after the completion of BCT. Although, the decreased leisure time physical activity prior to BCT might have predisposed Native Americans to a higher risk of injury during BCT which then carried over into AIT.

(b) In a study investigating risk-taking behaviors in Native Americans compared to Caucasians and African Americans, investigators found that male Native Americans reported higher rates of risk taking on 8 of 11 risk indicators.⁵⁰ This risk-taking behavior may manifest as a higher injury risk in AIT. Risk taking might be expected to increase traumatic injury rate more than overuse injuries, and, in fact, an increase in traumatic injury risk was seen in the present study. This may be at least a partial explanation for the increased injury risk in Native Americans when compared to Caucasians.

(2) Black men had higher injury risk for traumatic injuries compared to Caucasian men. In previous studies, Black men had a higher prevalence of lower extremity tendon injuries. In a study examining 865 U.S. military members who underwent Achilles tendon repair, the researchers found that Blacks had an overall increased risk of 4.2 (95 percent CI:3.6–4.7) for undergoing repair, when compared with non-Blacks.⁵¹ Another study (using data from 2000–2004 from the U.S. Defense Medical Epidemiology Database) found that, compared with Caucasian Service members, Black Service members had an adjusted rate ratio for quadriceps tendon tears of 2.9 (95 percent CI:2.4–3.4), patellar tendon tears of 4.5 (95 percent CI:3.9–5.2), and Achilles tendon tears 3.6 (95 percent CI:3.3–3.9).⁵² In a biomechanical study of the viscoelastic characteristics in the tricep surae between Black and Caucasian athletes, Black athletes were found to have a significantly greater muscle viscosity and muscle stiffness, which could result in tissue that is more likely to undergo failure if subject to sufficient trauma.⁵³ On the other hand, Blacks are less likely to experience stress fractures,

compared with Caucasians, possibly because Blacks having a higher bone density.⁵⁴⁻⁵⁶ Other studies performed during BCT and AIT have shown no differences when examining race and injury risk.^{6,10,24} The present study cannot determine why Black men reported more injuries than Caucasians, but the factors above may be worth exploring in future studies.

(3) Caucasian women had a higher risk of time-loss injuries compared to Black women. Other studies also suggest that Caucasian women have a higher risk of training injuries and stress fractures than other ethnic origins.^{10,54,57,58} In a study investigating knee-related disability among enlisted women in the U.S. Army, investigators found that non-Caucasian women had a 50 percent lower risk of knee-related discharge relative to Caucasian women.⁵⁹ In another study, investigators found that African American women had fewer blisters on their feet compared with Soldiers of other ethnicities.²⁶

d. Basic Training Site.

(1) For women, those arriving from Fort Leonard Wood had a higher injury rate when compared with those from Fort Jackson. The Service members arriving from Fort Jackson may have had a lower risk of being injured due to the multiple injury-reduction interventions introduced at that training facility.⁵ In 1998, the commander of the Fort Jackson Training Center increased the emphasis on reducing injury rates, and the USACHPPM established an injury coordinator position to provide state-of-the-art advice and material support to commanders and drill sergeants for reducing injury rates. Program monitoring from surveys and a Physical Training and Rehabilitation Program Surveillance System (a surveillance system developed by the Physical Therapy Department at Moncrief Army Community Hospital, Fort Jackson, to track injury information) suggests that these interventions were associated with a reduction in injury rates. Further, several other epidemiological consultations and studies have been performed at Fort Jackson, which may have raised awareness of injury prevention measures and subsequently reduced injury rates.^{8,11,13,16,20, 21,24}

(2) However, in the univariate analysis for time-loss injuries, men arriving from Fort Jackson had a higher injury rate when compared to those from Fort Knox. When examining previous self-reported injuries and tobacco use, there were no differences between Fort Jackson and Fort Knox. However, there was a small difference in muscular endurance and 2-mile run times with Fort Jackson having an approximate increase of 5 percent more Soldiers in the lowest quartiles for push-ups and sit-ups and slower 2-mile run times when compared to Fort Knox. This could be a partial reason for

the higher risk of injury for men arriving from Fort Jackson compared to those from Fort Knox.

e. Military Occupational Specialty. For men, injury risk was higher for the MOS of fuel and electrical systems repairer (63G) and the MOS of track vehicle mechanic (63Y) when compared to track vehicle repairer (63H). All three of these specialties have a “physical demands” rating of very heavy which means that on occasion men will have to lift over 100 pounds with frequent or constant lifting of in excess of 50 pounds. Also, all the men in these specialties perform repairs or maintenance on wheeled or tracked vehicles. When investigating current self-reported injury, tobacco use, and physical fitness scores among the three specialties, there were no differences. It is unclear why the MOS of fuel and electrical systems repairer and the MOS of track vehicle mechanic are at a higher risk of overuse injury when compared to the MOS of track vehicle repairer, when all three occupations have similar job objectives.

f. Self-Reported Injury.

(1) Injury risk in AIT was associated with current injuries perceived by the Soldier to have a negative effect on AIT performance. For men, injury risk for those with self-reported injuries in AIT was approximately 2.2 times higher for all injury categories (except traumatic injury (1.6)) and 1.5 times higher for women (except traumatic injury (2.2)). In answering this question, the Soldiers’ perception of injury limitations could have been influenced by the anticipation of the tasks they were required to perform as part of their MOS which can vary in terms of physical demands and duration of training. Therefore, if the Soldiers had an injury, it may or may not have affected their performance, but they could only answer this question as to how they perceived the level of difficulty associated with their MOS. Other factors that may have also influenced an individual’s perceived threat of injury could include age, gender, ethnicity, education, personality and peer pressure. Previous injury, both overuse and traumatic, have also been shown to place Soldiers at a higher risk of re-injury^{10,60,61} In one study, having a traumatic injury increased the risk of subsequent injury by 83 percent when compared to having an injury categorized as an overuse injury or unspecified pain.⁶²

(2) In civilian studies, investigators have shown that those who sustained an injury (both traumatic and overuse) within the last year, as well as those who sustained a current injury, had a higher risk of re-injury than those who reported no previous or current injuries.^{38,63-67} In a study investigating low back pain as a risk factor for recurrent injuries in varsity athletes, researchers found that athletes who reported a previous low back injury were at a three times greater risk and athletes who reported current back pain were at a six times greater risk of sustaining a lower back injury.⁶⁷ It

is possible that previous or current injuries may not be the cause of subsequent injuries if the former are treated adequately. In a study investigating re-injury rates of amateur male soccer players, researchers found an 11 percent re-injury rate for a coach-controlled rehabilitation group (coaches received a 10-step rehabilitation program to implement as a guide for functional rehabilitation after an injury occurred) compared to a 29 percent re-injury rate for the control group (coaches were instructed to go on with training and management of injuries as usual).⁶⁸

(3) In the present study, Soldiers were asked to report on injuries believed to affect AIT training. Soldiers who answered affirmatively may or may not have sought out medical attention for those specific injuries, or the injury may have increased susceptibility to other injuries. Other studies have demonstrated that inadequate rehabilitation and a premature return to competition are risk factors for reinjury.^{31,69,70}

g. Cigarette Use.

(1) For men, injury risk was higher in smokers than nonsmokers and demonstrated a dose-response relationship in which injury risk increases with the number of cigarettes smoked per day. Previous studies have also demonstrated an increased risk of injury in smokers compared to nonsmokers, the number of cigarettes smoked per day, and risk of musculoskeletal injury.^{5,13,35,46-48,71-74} The relationship between tobacco use and injury may be due to a compromised ability to repair damaged tissues, thereby increasing susceptibility to the repetitive microtrauma that presumably causes overuse injuries.⁷⁵ In a study examining the healing of tibial fractures, investigators found that healing time to clinical union was 166 ± 92 days for smokers versus 134 ± 71 days for nonsmokers (a 24 percent slower healing time). Possible attributes of the delayed time to clinical union could be decreased oxygen saturation levels and/or impaired blood flow to the injured area in smokers.⁷⁶ In examining the healing of wounds, researchers found that 5 out of 15 patients who underwent intraoral bone grafting with simultaneous implant placement experienced impaired wound healing. Of these five, four admitted to smoking in the preoperative period. Vasoconstriction of the tissues due to nicotine was one of the suggested causes of the delayed healing.⁷⁷ Therefore, the constant physical stressors of AIT training may result in weaken tissues from training and overuse, which may cause a greater susceptibility of injury in smokers.

(2) For traumatic injuries, the relationship between tobacco use and musculoskeletal injury may be due to greater risk-taking behaviors. In an Air Force study, recruits who were cigarette smokers had higher scores than nonsmokers on various measures of risk taking. These included an overall measure of risk-taking, in

addition to greater rebelliousness, less seat belt use, more risky sex, a more favorable view of illegal drug use, more alcohol use, more binge drinking, less physical activity, less intake of fruits and vegetables, and greater intake of high fat foods.³²

(a) In civilian studies, smokers had more motor vehicle accidents, more traffic violations, less seat belt use, less physical activity, more alcohol consumption and lower intake of fruits and vegetables compared to nonsmokers.⁷⁸⁻⁸⁰

(b) Another hypothesis to explain the association between injuries and tobacco use may be due to a decrease in fatigue resistance. In a study investigating musculoskeletal fatigue and smoking history, investigators found that the skeletal muscle of smokers was more fatigable than that of older and physically active matched nonsmokers.⁸¹ The authors suggested that smoking may have an acute and reversible effect on skeletal muscle fatigability caused by carbon monoxide in cigarette smoke. Carbon monoxide can reduce the oxygen content of the blood by binding to the oxygen sites of the hemoglobin which then diminishes oxygen delivery and results in oxygen being released at a slower rate to the tissues.⁸² In smokers, carboxyhemoglobin may reach levels of 9 percent.⁸²

(c) Other investigators have found that fatigue leads to decrements in proprioceptive ability, a decrease in joint stability, and alterations in muscle activity which can then possibly lead to a higher risk of injury.^{83,84}

h. Smokeless Tobacco Use.

(1) For men, risk of injury among frequent smokeless tobacco users was higher than among non-users. In a study investigating injury proneness in infantry conscripts, investigators found that smokeless tobacco placed conscripts at a 2.4 times higher risk of injury than non-smokeless users.³⁵ In another study investigating tobacco use and injury risk among military parachutists, investigators found that smokeless tobacco users had a 50 percent greater odds of injury than non-smokeless users (although the association was not statistically significant at the .05 level).⁸⁵ It has been demonstrated that concentration levels of nicotine and cotinine in the blood are similar in cigarette and smokeless tobacco users.^{86,87} Therefore, any harmful effects of nicotine exposure related to cigarettes would also be an expected hazard of smokeless tobacco.

(2) As noted earlier, one of the effects of nicotine is vasoconstriction of the tissues which has been suggested to lead to delayed healing.⁷⁷ When examining the effects of smokeless tobacco on the cardiorespiratory response to submaximal exercise, investigators have found an increase in heart rate as well as a decrease in

stroke volume after their subjects used smokeless tobacco compared to a placebo group.⁸⁸ Investigators also found increased plasma lactate concentration levels at any given submaximal oxygen uptake after their subjects used smokeless tobacco, which is suggestive of a greater demand for glycolytic energy production.⁸⁸ Faster heart rates and greater use of glycolytic energy sources during submaximal exercise could then lead to fatigue and a higher risk of injury. However, other studies have shown that nicotine does not influence the perception of exertion at low intensity exercise, that long-term use of smokeless tobacco does not influence exercise capacity in healthy and physically well-trained subjects, and that smokeless tobacco has no effect on reaction time.⁸⁹⁻⁹¹ It is possible that the increased risk of injury associated with smokeless tobacco use could be attributed to a number of different conditions.

(3) In summary, long-term effects of smokeless tobacco could lead to slower healing times, with acute effects to include fatigue due to an increased heart rate, a greater dependence on blood glucose as fuel during rest and sub-maximal exercise, and a delay in the nervous transmission across the neuromuscular junction.^{88,92,93}

i. Muscular Endurance. Time-loss injury risk was higher for Soldiers who had poor muscular endurance (push-ups and sit-ups), which has also been shown to be associated with injuries in BCT.^{10,11,13,24,72,94,95} Many tasks performed in BCT and AIT require muscular endurance of the upper body. Lack of muscular endurance could lead to fatigue and a greater reliance on different muscle groups as the active muscles begin to fatigue.^{96,97} This unaccustomed stress may increase the risk of injury.

j. Two-Mile Run Times. Injury risk for the slowest 2-mile run times was higher for men and women when compared to the fastest 2-mile run times. Previous studies investigating run times during BCT have also found that slower run times place Soldiers at a higher risk of injury.^{4,11,13,17,98} The Soldiers with the slowest 2-mile run times would have lower aerobic capacities than those with the fastest 2-mile run times.⁹⁹ Soldiers with lower aerobic capacities will likely experience greater physiological stress and/or fatigue during AIT tasks (such as running, cross-training and calisthenics) due to exercising at a higher percentage of their maximum aerobic capacity when compared to Soldiers with greater fitness levels. Soldiers of lower fitness levels will not only be exercising at a higher percentage of their aerobic capacity to accomplish the same task as a more fit Soldier, but they will also perceive tasks as more difficult.¹⁰⁰ The greater physiological stress and/or fatigue experienced may lead to a higher risk of injury. Studies on fatigue have demonstrated decrements in proprioceptive ability; a decrease in joint stability; alterations in muscle activity; and changes in gait, balance, low frequency fatigue, neuromuscular function, and ligament laxity.^{83,84,101-110}

9. CONCLUSIONS. This study identified risk factors for time-loss injury in Ordnance School AIT Soldiers. Overall, 31 percent of men and 54 percent of women involved in the project incurred at least one time-loss injury. When examining injury risk for all four injury categories, both cigarette use and self-reported injury were associated with a higher risk of injury in men. For three out of the four injury categories, self-reported injury (women), sit-ups (men), and the 2-mile run (men and women) were associated with a higher risk of injury.

10. RECOMMENDATIONS. In an effort to reduce injuries, surveillance and tracking of injuries in AIT Soldiers could alert commanders to elevated levels of injuries or to injury outbreaks. Smoking cessation classes and fitness training prior to entry are potential strategies to reduce injuries.

11. TECHNICAL ASSISTANCE. Direct inquiries regarding this report to Mr. Tyson Grier, Project Officer, Injury Prevention Program, Directorate of Epidemiology and Disease Surveillance, at commercial (410) 436-5450, DSN 584-5450, or email to tyson.grier@us.army.mil.

TYSON L. GRIER
Health Scientist

Approved:

BRUCE H. JONES
Program Manager, Injury Prevention

APPENDIX A

REFERENCES

1. Army (2005) Army Regulation 40-5: Preventive Medicine. Washington DC: Headquarters, Department of the Army.
2. Defense Safety Oversight Council (2006) DoD Military Injury Prevention Priorities Working Group: Leading Injuries, Causes and Mitigation Recommendations. Washington, D.C. Defense Safety Oversight Council.
3. Almeida S, Williams M, Shaffer R, Luz J, Badong K, Brodine S (1997) A physical training program to reduce musculoskeletal injuries in U.S. Marine Corps recruits. Naval Health Research Center, Technical Report No. NHRC07-2B, National Technical Information Service, U.S. Dept of Commerce
4. Knapik JJ, Ang P, Reynolds K, Jones B. Physical fitness, age and injury incidence in infantry soldiers. *Journal of Occupational Medicine*, 1993; 35: 598-603.
5. Primary Prevention of Injuries in Initial Entry Training. Knapik JJ, Hauret KG, Jones BH. (2006) Textbook of Military Medicine. Recruit Medicine. Lenhart MK, Lounsbury DE, North RB. Washington DC: Bordon Institute.
6. Henderson NE, Knapik JJ, Shaffer SW, McKenzie TH, Schneider GM. Injuries and injury risk factors among men and women in US Army combat medic advanced individual training. *Military Medicine*, 2000; 165: 647-652.
7. Grier T, Knapik J, Canada S, Canham-Chervak M, Jones B. Risk factors for time-loss injuries in the United States Army Ordnance School Advanced Individual Training. *Medicine and Science in Sports and Exercise*, 2009; 41: S574.
8. Knapik JJ, Cuthie J, Canham M, Hewitson W, Laurin MJ, Nee MA, Hoedebecke E, Hauret K, Carroll D, Jones BH (1998) Injury incidence, injury risk factors, and physical fitness of U.S. Army basic trainees at Ft Jackson SC, 1997. 29-HE-7513-98, Aberdeen Proving Ground, MD: U.S. Army Center for Health Promotion and Preventive Medicine.
9. Associations among body composition, physical fitness, and injuries in men and women Army trainees Jones BH, Bovee MW, Knapik JJ. (1992) Body Composition and Physical Performance. Marriott BM, Grumstrup-Scott J. Washington, D.C.: National Academy Press, 141-173.
10. Jones BH, Cowan DN, Tomlinson JP, Robinson JR, Polly DW, Frykman PN. Epidemiology of injuries associated with physical training among young men in the Army. *Medicine and Science in Sports and Exercise*, 1993; 25: 197-203.
11. Jones BH, Bovee MW, Harris JM, Cowan DN. Intrinsic risk factors for exercise-related injuries among male and female Army trainees. *American Journal of Sports Medicine*, 1993; 21: 705-710.

12. Bensel CK, Kish RN (1983) Lower extremity disorders among men and women in Army basic training and effects of two types of boots. TR-83/026, Natick, MA: U.S. Army Natick Research and Development Laboratories
13. Knapik JJ, Sharp MA, Canham-Chervak M, Hauret K, Patton JF, Jones BH. Risk factors for training-related injuries among men and women in Basic Combat Training. *Medicine and Science in Sports and Exercise*, 2001; 33: 946-954.
14. Knapik JJ, Hauret KG, Arnold S, Canham-Chervak M, Mansfield AJ, Hoedebecke EL, McMillian D. Injury and fitness outcomes during implementation of Physical Readiness Training. *International Journal of Sports Medicine*, 2003; 24: 372-381.
15. Knapik JJ, Darakjy S, Scott S, Hauret KG, Canada S, Marin R, Palkoska F, VanCamp S, Piskator E, Rieger W, Jones BH (2004) Evaluation of two Army fitness programs: the TRADOC Standardized Physical Training Program for Basic Combat Training and the Fitness Assessment Program. 12-HF-5772B-04, Aberdeen Proving Ground, MD: US Army Center for Health Promotion and Preventive Medicine.
16. Westphal KA, Friedl KE, Sharp MA, King N, Kramer TR, Reynolds KL, Marchitelli LJ (1995.) Health, performance and nutritional status of U.S. Army women during basic combat training. T96-2, Natick, MA: U.S. Army Research Institute of Environmental Medicine.
17. Knapik J, Swedler D, Grier T, Hauret K, Bullock S, Williams K, Darakjy S, Lester M, Tobler S, Clemmons N, Jones B (2008) Injury reduction effectiveness of prescribing running shoes based on foot shape in basic combat training. 12-MA-05SB-08, Aberdeen Proving Ground, MD US Army Center for Health Promotion and Preventive Medicine
18. Bell NS, Mangione TW, Hemenway D, Amoroso PJ, Jones BH. High injury rates among female Army trainees. A function of gender? *American Journal of Preventive Medicine*, 2000; 18(Suppl. 3): 141-146.
19. Jones BH. Injuries among men and women in gender-integrated BCT units. Ft Leonard Wood 1995. *Medical Surveillance Monthly Report*, 1996; 2: 2-3,7-8.
20. Canham-Chervak M, Knapik JJ, Hauret K, Cuthie J, Craig S, Hoedebecke E (2000.) Determining physical fitness entry criteria for entry into Army Basic Combat Training: can these criteria be based on injury? 29-HE-1395-00, Aberdeen Proving Ground, MD: US Army Center for Health Promotion and Preventive Medicine.
21. Knapik JJ, Hauret K, Bednarek JM, Arnold S, Canham-Chervak M, Mansfield A, Hoedebecke E, Mancuso J, Barker TL, Duplessis D, Heckel H, Peterson J, (2001) TSotAPFS (2001) The Victory Fitness Program: Influence of the US Army's emerging physical fitness doctrine on fitness and injuries in Basic Combat Training. 12-MA-5762-01, Aberdeen Proving Ground, MD US Army Center for Health Promotion and Preventive Medicine.

22. Knapik JJ, Darakjy S, Scott SJ, Hauret KG, Canada S, Marin R, Rieger W, Jones BH. Evaluation of a standardized physical training program for Basic Combat Training. *Journal of Strength and Conditioning Research*, 2005; 19: 246-253.
23. Kowal DM. Nature and causes of injuries in women resulting from an endurance training program. *American Journal of Sports Medicine*, 1980; 8: 265-269.
24. Knapik JJ, Sharp MA, Canham ML, Hauret K, Cuthie J, Hewitson W, Hoedebecke E, Laurin MJ, Polyak C, Carroll D, Jones B (1999.) Injury incidence and injury risk factors among US Army Basic Trainees at Ft Jackson, SC. 29-HE-8370-99, Aberdeen Proving Ground MD US Army Center for Health Promotion and Preventive Medicine.
25. Jones B, Shaffer R, Snedecor M. Injuries treated in outpatient clinics: surveys and research data. *Military Medicine*, 1999; 164(suppl): 6-1-6-89.
26. Knapik JJ, Reynolds KL, Barson J (1997) Influence of antiperspirants on foot blisters following road marching. ARL-TR-1333, Aberdeen Proving Ground, MD U.S. Army Research Laboratory
27. Knapik JJ, Canham-Chervak M, Hauret K, Laurin MJ, Hoedebecke E, Craig S, Montain S. Seasonal variations in injury rates during US Army Basic Combat Training. *Annals of Occupational Hygiene*, 2002; 46: 15-23.
28. Rehabilitation and Reconditioning Potach D, Borden R. (2000) Essentials of strength training and conditioning. Baechle T. Hong Kong: Human Kinetics.
29. Diagnosing and treating your sports injury Micheli L, Jenkins M. (2001) The sports medicine bible. HarperCollins. New York: HarperCollins.
30. Brown M, Brown R. Athletic Injuries. *Trauma*, 1999; 1: 271-278.
31. McKay GD, Goldie PA, Oakes BW. Ankle injuries in basketball: injury rate and risk factors. *British Journal of Sports Medicine*, 2001; 35:
32. Lando H, Haddock C, Klesges R, Talcott G, Jensen J. Smokeless tobacco use in a population of young adults. *Addictive Behaviors*, 1999; 24: 431-437.
33. Croisier J. Muscular imbalance and acute lower extremity muscle injuries in sport. *International Journal of Sports Medicine*, 2004; 5: 169-176.
34. Knapik JJ, Bauman CL, Jones BH, Harris JM, Vaughan L. Preseason strength and flexibility imbalances associated with athletic injuries in female collegiate athletes. *American Journal of Sports Medicine*, 1991; 19: 76-81.
35. Heir T, Eide G. Injury proneness in infantry conscripts undergoing a physical training programme: smokeless tobacco use, higher age, and low levels of physical fitness are risk factors. *Scandinavian Journal of Medicine and Science in Sports*, 1997; 7: 304-311.
36. Barowclough F. The process of aging. *Journal of Advanced Nursing*, 1981; 6: 319-325.
37. Macera CA, Jackson KL, Hagenmaier GW, Kronenfeld JJ, Kohl HW, Blair SN. Age, physical activity, physical fitness, body composition and incidence of orthopedic problems. *Research Quarterly for Exercise and Sport*, 1989; 60: 225-233.

38. Macera CA, Pate RR, Powell KE, Jackson KL, Kendrick JS, Craven TE. Predicting lower-extremity injuries among habitual runners. *Archives of Internal Medicine*, 1989; 49: 2565-2568.
39. Colbert LH, Hootman JM, Macera CA. Physical activity-related injuries in walkers and runners in the Aerobics Center Longitudinal Study. *Clinical Journal of Sport Medicine*, 2000; 10: 259-263.
40. Hootman JM, Macera CA, Ainsworth BA, Martin M, Blair SN. Predictors of lower extremity injury among recreationally active adults. *Clinical Journal of Sport Medicine*, 2002; 12: 99-106.
41. Taunton JE, Ryan MB, Clement DB, McKenzie DC, Lloyd-Smith DR, Zumbo BD. A prospective study of running injuries: the Vancouver Sun Run "In Training" clinics. *British Journal of Sports Medicine*, 2003; 37: 239-244.
42. Orchard JW. Intrinsic and extrinsic risk factors for muscle strains in Australian football. *American Journal of Sports Medicine*, 2001; 29: 300-303.
43. Bennell KL, Crossley K. Musculoskeletal injuries in track and field: incidence, distribution and risk factors. *Australian Journal of Science and Medicine in Sport*, 1996; 28: 69-75.
44. CDC. Cigarette smoking among adults--United States, 2000. *MMWR*, 2002; 51: 642-645.
45. Clark D, Holtzman D, Cobb N. Surveillance for health behaviors of american Indians and alaska natives, 1997-2000. *MMWR Surveillance Summaries*, 2003; 52: 1-13.
46. Altarac M, Gardner JW, Popovich RM, Potter R, Knapik JJ, Jones BH. Cigarette smoking and exercise-related injuries among young men and women. *American Journal of Preventive Medicine*, 2000; 18 (Suppl 3S): 96-102.
47. Munnoch K, Bridger R. Smoking and injury in Royal Marines training. *Occupational Medicine*, 2007; 57: 214-216.
48. Reynolds KL, Heckel HA, Witt CE, Martin JW, Pollard JA, Knapik JJ, Jones BH. Cigarette smoking, physical fitness, and injuries in infantry soldiers. *American Journal of Preventive Medicine*, 1994; 10: 145-150.
49. Health and Human Services (2000). Healthy people 2010 2nd ed. Atlanta, GA: Center for Disease Control and Prevention, 2000.
50. Gruber E, Dicemente R. Risk-taking behavior among native american adolescents in Minnesota public schools: comparisons with black and white adolescents. *Ethnicity and Health*, 1996; 1: 261-267.
51. Davis J, Mason K, Clark D. Achilles tendon ruptures stratified by age, race, and cause of injury among active duty U.S. military members. *Military Medicine*, 1999; 164: 872-873.
52. Owens B, Mountcastle S, White D. Racial differences in tendon rupture incidence. *International Journal of Sports Medicine*, 2007; 28: 617-620.

53. Fukashiro S, Abe T, Shibayama A, Brechue W. Comparison of viscoelastic characteristics in triceps surae between black and white athletes. *Aca Physiol Scand*, 2002; 175: 183-187.
54. Brudvig TGS, Gudger TD, Obermeyer L. Stress fractures in 295 trainees: a one-year study of incidence as related to age, sex, and race. *Military Medicine*, 1983; 148: 666-667.
55. Trotter M, Broman GE, Peterson RR. Densities of white and negro skeleton. *Journal of Bone and Joint Surgery*, 1960; 42A: 50-58.
56. Barondess D, Nelson D, Schlaen S. Whole body bone, fat, and lean mass in black and white men. *Journal of Bone and Mineral Research*, 1997; 12: 967-971.
57. Gardner LI, Dziados JE, Jones BH, Brundage JF, Harris JM, Sullivan R, Gill P. Prevention of lower extremity stress fractures: a controlled trial of a shock absorbent insole. *American Journal of Public Health*, 1988; 78: 1563-1567.
58. Knapik JJ, Jones BH, Hauret KG (2003). Injury control in the United States military.
59. Sulsky S, Mundt K, Bigelow C, Amoroso P. Risk factors for occupational knee related disability among enlisted women in the US Army. *Occupational Environmental Medicine*, 2002; 59: 601-607.
60. Knapik JJ, Jones SB, Darakjy S, Hauret KG, Nevin R, Grier T, Jones BH. Injuries and injury risk factors among members of the United States Army Band. . *American Journal of Industrial Medicine*, 2007; 50: 951-961.
61. Bender JA, Pierson JK, Kaplan HM, Johnson AJ. Factors affecting the occurrence of knee injuries. *Journal of the Association for Physical and Mental Rehabilitation*, 1964; 18: 130-134.
62. Schneider GA, Bigelow C, Amoroso PJ. Evaluating risk of re-injury among 1214 Army Airborne soldiers using a stratified survival model. *American Journal of Preventive Medicine*, 2000; 18(Suppl3): 156-163.
63. Emery C, Meeuwisse W, Hartmann S. Evaluation of risk factors for injury in adolescent soccer; implementation and validation of an injury surveillance system. *The American Journal of Sports Medicine*, 2005; 33: 1882-1891.
64. Taimela S, Kujala UM, Newman K. Intrinsic risk factors and athletic injuries. *Sports Medicine*, 1990; 9: 205-215.
65. Madden C, Garrett J, Cole T, Runge J, Porter C. The urban epidemiology of recurrent injury: beyond age, race, and gender stereotypes. *Academic Emergency Medicine*, 1997; 4: 772-775.
66. Walters SD, Hart LE, McIntosh JM, Sutton JR. The Ontario cohort study of running-related injuries. *Archives of Internal Medicine*, 1989; 149: 2561-2564.
67. Greene H, Cholewicki J, Galloway M, Nguyen C, Radebold A. A history of low back injury is a risk factor for recurrent back injuries in varsity athletes. *The American Journal of Sports Medicine*, 2001; 29: 795-800.

68. Hagglund M, Walden M, Ekstrand J. Lower reinjury rate with a coach-controlled rehabilitation program in amateur male soccer; a randomized controlled trial. *The American Journal of Sports Medicine*, 2007; 35: 1433-1442.
69. Murphy D, Connolly D, Beynnon B. Risk factors for lower extremity injury: a review of the literature. *British Journal of Sports Medicine*, 2003; 37: 13-29.
70. Ekstrand J, Gillquist J. The avoidability of soccer injuries. *International Journal of Sports Medicine*, 1983; 4: 124-128.
71. Reynolds KL, Heckel HA, Martin JW, Pollard JA, Knapik JJ, Jones BH. Cigarette smoking, physical fitness, and injuries in infantry soldiers. *Am J Prev Med*, 1994; 10: 145-150.
72. Knapik J, Swedler D, Grier T, Hauret K, Bullock S, Williams K, Darakjy S, Lester M, Tobler S, Jones B. Injury reduction effectiveness of selecting running shoes based on plantar shape. *Journal of Strength and Conditioning Research*, 2009; 23: 685-697.
73. Dettori J, Reynolds K, Amoroso P, Barnes J, Westphal K, Lavin P (1996). Smoking and injury risk among female US army basic combat trainees. Presented at the Third International Conference for Injury Prevention and Control. Melbourne, Australia
74. Reynolds K, Amoroso P, Dettori J. Association of tobacco use with injuries among infantry soldiers carrying loads on a 100 mile road march. *Presented at the Third International Conference for Injury Prevention and Control*, 1996;
75. Amoroso PJ, Reynolds KL, Barnes JA, White DJ (1996) Tobacco and injuries: an annotated bibliography. TN96-1, Natick, MA: U.S. Army Research Institute of Environmental Medicine.
76. Kyro A, Usenius J, Aarnio M, Kunnamo I, Avidainen V. Are smokers a risk group for delayed healing of tibial shaft fractures. *Annales Chirurgiae et Gynaecologiae*, 1993; 82: 254-262.
77. Jones JK, Triplett RG. The relationship of cigarette smoking to impaired intraoral wound healing. *Journal of Oral and Maxillofacial Surgery*, 1992; 50: 237-239.
78. Difranza J, Winters T, Goldberg R, Cirillo L, Biliouris T. The relationship of smoking to motor vehicle accidents and traffic violations. *New York State Journal of Medicine*, 1986; 86: 464-467.
79. Eiser J, Sutton S, Wober M. Smoking, seat belts, and beliefs about health. *Addictive Behaviors*, 1979; 4: 331-338.
80. Chiolero A, Wietlisbach V, Ruffieux C, Paccaud F, Cornuz J. Clustering of risk behaviors with cigarette consumption: a population-based survey. *Preventive Medicine*, 2006; 42: 348-353.
81. Wust RC, Morse CI, DeHaanJornRittweger A, Jones DA, Degens H. Skeletal muscle properties and fatigue resistance in relation to smoking history. *European Journal of Applied Physiology*, 2008; 104: 103-110.


82. Rietbrock N, Kunkel S, Worner W, Eyer P. Oxygen-dissociation kinetics in the blood of smokers and non-smokers: interaction between oxygen and carbon monoxide at the hemoglobin molecule. *Archives of Pharmacology*, 1992; 345: 123-128.
83. Rozzi S, Lephart S, Fu F. Effects of muscular fatigue on knee joint laxity and neuromuscular characteristics of male and female athletes. *Journal of Athletic Training*, 1999; 34: 106-114.
84. Melnyk M, Gollhofer A. Submaximal fatigue of the hamstrings impairs specific reflex components and knee stability. *Knee surgery, sports traumatology, arthroscopy*, 2007; 15: 525-532.
85. Amoroso P, Dettori J, Reynolds K. Tobacco use and injury risk among military parachutists. *Third International Conference for Injury Prevention and Control*. Melbourne, Australia, 1996
86. Benowitz NL, Porchet H, Sheiner L, Jacobs P. Nicotine absorption and cardiovascular effects with smokeless tobacco use: Comparison with cigarettes and nicotine gum. *Clinical Pharmacology and Therapeutics*, 1988; 44: 23-28.
87. Gritz E, Baer-Weiss V, Benowitz N, Vunakis HV, Jarvik M. Plasma nicotine and cotinine concentrations in habitual smokeless tobacco users. *Clinical Pharmacology and Therapeutics*, 1981; 30: 201-209.
88. Duser BV, Raven P. The effects of oral smokeless tobacco on the cardiorespiratory response to exercise. *Medicine and Science in Sports and Exercise*, 1992; 24: 389-395.
89. Perkins K, Sexton J, Solberg-Kassel R, Epstein L. Effects of nicotine on perceived exertion during low-intensity activity. *Medicine and Science in Sports and Exercise*, 1991; 23: 1283-1288.
90. Bolinder G, Noren A, Faire WD, Faire UD. Long-term use of smokeless tobacco and physical performance in middle-aged men. *European Journal of Clinical Investigation*, 1997; 27: 427-433.
91. Escher S, Tucker A, Lundin T, Grabiner M. Smokeless tobacco, reaction time, and strength in athletes. *Medicine and Science in Sports and Exercise*, 1998; 30: 1548-1551.
92. Colberg S, Casazza G, Horning M, Brooks G. Increased dependence on blood glucose in smokers during rest and sustained exercise. *Journal of Applied Physiology*, 1994; 76: 26-32.
93. Lester M, Jacobson B, Edwards S. Effect of ingestion of smokeless tobacco on motor distal latency of the median nerve. *Perceptual and Motor Skills*, 1988; 67: 283-290.
94. Jones BH, Manikowski R, Harris JR, Dziados J, Norton S, Ewart T, Vogel JA (1988) Incidence of and risk factors for injury and illness among male and female Army basic trainees. T19/88, Natick, MA: United States Army Research Institute of Environmental Medicine.

95. Knapik JJ, Bullock SH, Canada S, Toney E, Wells JD, Hoedebecke E, Hauret KG, Rieger W, Palkoska F, VanCamp S, McMillian D, Edwards D, Billet M (2003) The Aberdeen Proving Ground Injury Control Project: Influence of a multiple intervention program on injuries and fitness among Ordnance School students in Advanced Individual Training. 12-HF-7990-03, Aberdeen Proving Ground, MD US Army Center for Health Promotion and Preventive Medicine
96. Gleeson N, Reilly T, Mercer T, Rakowski S, Rees D. Influence of acute endurance activity on leg neuromuscular and musculoskeletal performance. *Medicine and Science in Sports and Exercise*, 1998; 30: 596-608.
97. Hirsohi A, Foley J, Prior B, Dudley G, Meyer B. Vastus lateralis fatigue alters recruitment of musculus quadriceps femoris in humans. *Journal of Applied Physiology*, 2002; 92: 679-684.
98. Hauret KG, Knapik JJ, Darakjy S, Canada S, Marin RE, Jones BH. Reduced injury risk in Army basic combat training with a standardized physical training program. *Medicine and Science in Sports and Exercise*, 2004; 36: S309.
99. Knapik JJ. The Army Physical Fitness Test (APFT): a review of the literature. *Military Medicine*, 1989; 154: 326-329.
100. Garcin M, Vautier JF, Vandewalle H, Monod H. Rating of perceived exertion (RPE) as an index of aerobic endurance during local and general exercise. *Ergonomics*, 1988; 41: 105-114.
101. Candau R, Belli A, Millet GY, George D, Barbier B, Rouillon JD. Energy cost and running mechanics during a treadmill run to voluntary exhaustion in humans. *European Journal of Applied Physiology*, 1998; 77: 479-485.
102. Nyland JA, Shapiro R, Stine RL, Horn TS, Ireland ML. Relationship of fatigued run and rapid stop to ground reaction forces, lower extremity kinematics, and muscle activation. *Journal of Orthopedic and Sports Physical Therapy*, 1994; 20: 132-137.
103. Gerlach K, White S, Burton H, Dorn J, Leddy J, Horvath P. Kinetic changes with fatigue and relationship to injury in female runners. *Medicine and Science in Sports and Exercise*, 2005; 37: 657-663.
104. LeBris R, Billat V, Auvinet B, Chaleil D, Hammard L, Barrey E. Effect of fatigue on stride pattern continuously measured by an accelerometric gait recorder in middle distance runners. *Journal of Sports Medicine and Physical Fitness*, 2006; 46: 227-231.
105. Yoshino K, Motoshige T, Araki T, Matsuoka K. Effect of prolonged free-walking fatigue on gait and physiological rhythm. *Journal of Biomechanics*, 2004; 37: 1271-1280.
106. Johnston R, Howard M, Cawley P, Losse G. Effect of lower extremity muscular fatigue on motor control performance. *Medicine and Science in Sports and Exercise*, 1998; 30: 1703-1707.

107. Dickin D, Doan J. Postural stability in altered and unaltered sensory environments following fatiguing exercise of lower extremity joints. *Scandinavian Journal of Medicine and Science in Sports*, 2008; 18: 765-772.
108. Baptista R, Scheeren E, Macintosh B, Vaz M. Low-frequency fatigue at maximal and submaximal muscle contractions. *Brazilian Journal of Medical and Biological Research*, 2009; 42: 380-385.
109. Wojtys E, Wylie B, Huston L. The effects of muscle fatigue on neuromuscular function and anterior tibial translation in healthy knees. *American Journal of Sports Medicine*, 1996; 24: 615-621.
110. Sakai H, Tanaka S, Kurosawa H, Masujima A. The effect of exercise on anterior knee laxity in female basketball players. *International Journal of Sports Medicine*, 1992; 13: 552-554.

APPENDIX B

QUESTIONNAIRE FOR ORDNANCE SCHOOL SOLDIERS (EXAMPLE)

 58295

Soldier Health Inprocessing Sheet, revised 7Jun01
ALL SOLDIERS FILL OUT THE FOLLOWING

1. Today's Date (DD-MM-YY) - -

3. Last Name

4. First Name

5. Grade or
Enlisted Officer

6. Race
☐ Asian American ☐ Hispanic American
☐ African American ☐ Native American
☐ Caucasian American ☐ Other American

7. Gender
☐ Male ☐ Female

8. Date of Birth (DD-MM-YY) - -

9. Age

10. Unit Assigned to:
☐ A (16th) ☐ B (16th) ☐ C (16th) ☐ D (16th) ☐ E (16th) ☐ A (143rd) ☐ B (143rd) ☐ C (143rd)

11. Basic Training Site:
☐ Ft. Jackson ☐ Ft. Knox ☐ Ft. Leonard Wood ☐ Ft. Benning ☐ Ft. Sill ☐ Other

12. Do you presently have an injury that would adversely affect your performance during AIT?
☐ Yes ☐ No

13. Do you presently have an illness that would adversely affect your performance during AIT?
☐ Yes ☐ No

14. If your answer to Question #12 or #13 is Yes, what area of the body does the injury or illness affect?
☐ General Health ☐ Arm ☐ Lower Back ☐ Ankle
☐ Eyes ☐ Hand ☐ Hip and Upper Leg ☐ Foot
☐ Head ☐ Neck and Upper Back ☐ Knee ☐ Other
☐ Shoulder ☐ Chest ☐ Lower Leg

15. When were you injured?
☐ Prior to BCT ☐ During BCT ☐ After BCT

16. When did your illness begin?
☐ Prior to BCT ☐ During BCT ☐ After BCT

17. In the space provided, tell us why you may need to see the doctor:

18. Did you smoke 1 or more cigarettes in the 30 days before Basic Training? ☐ Yes ☐ No

19. Did you smoke on 20 or more days in the 30 days before Basic Training? ☐ Yes ☐ No
If yes, how many cigarettes? ☐ 10 or fewer cigarettes per day on average
☐ 10-20 cigarettes per day on average
☐ 20 or more cigarettes per day on average

20. Did you use smokeless tobacco (chewing, snuffing, pinching, etc.) at least once in the 30 days before Basic Training? ☐ Yes ☐ No



21. Did you use smokeless tobacco (chewing, snuffing, pinching, etc.) on 20 or more days in the 30 days before Basic Training? ☐ Yes ☐ No
If yes, how much? ☐ Less than 1 can, pouch, or plug per day on average
☐ 1 can, pouch, or plug per day on average
☐ 2 or more cans, pouches, or plugs per day on average

FEMALES ONLY:

22. Have you had a PAP smear in the last year? ☐ Yes ☐ No
If yes, were the results abnormal? ☐ Yes ☐ No

APPENDIX C

INJURY SHEET FOR CLINICIANS (EXAMPLE)

  Draft	Injury Sheet revised June 2001	Today's date (DD/MM/YY)
Last Name <div style="border: 1px solid black; width: 300px; height: 1.2em; margin: 2px 0;"></div>		<div style="border: 1px solid black; width: 40px; height: 1.2em; margin: 2px 0;"></div> - <div style="border: 1px solid black; width: 40px; height: 1.2em; margin: 2px 0;"></div> - <div style="border: 1px solid black; width: 40px; height: 1.2em; margin: 2px 0;"></div>
Gender: <input type="checkbox"/> Male <input type="checkbox"/> Female	Unit: <input type="checkbox"/> A (16th) <input type="checkbox"/> B (16th) <input type="checkbox"/> C (16th) <input type="checkbox"/> D (16th) <input type="checkbox"/> E (16th) <input type="checkbox"/> A (143rd) <input type="checkbox"/> B (143rd) <input type="checkbox"/> C (143rd) <input type="checkbox"/> Permanent Party <input type="checkbox"/> USMC <input type="checkbox"/> USAF <input type="checkbox"/> ANCOG <input type="checkbox"/> BNCOC <input type="checkbox"/> Other	

1. Cause Codes (check one) <input type="checkbox"/> Sports <input type="checkbox"/> Running <input type="checkbox"/> PT (other than running) <input type="checkbox"/> Road March <input type="checkbox"/> Environmental (heat, cold) <input type="checkbox"/> Fall <input type="checkbox"/> Work Related <input type="checkbox"/> Fighting, anger-related <input type="checkbox"/> Other <div style="border: 1px solid black; width: 100px; height: 1.2em; display: inline-block;"></div> <input type="checkbox"/> Unknown 2. Location (check one) <input type="checkbox"/> Left <input type="checkbox"/> Right <input type="checkbox"/> Bilateral <input type="checkbox"/> Other <input type="checkbox"/> Unknown 3. Body Part (check one) <input type="checkbox"/> Unknown <input type="checkbox"/> Other Head <input type="checkbox"/> Ear <input type="checkbox"/> Eye <input type="checkbox"/> Nose <input type="checkbox"/> Neck <input type="checkbox"/> Face, NOS <input type="checkbox"/> Head, NOS Shoulder <input type="checkbox"/> Clavicle <input type="checkbox"/> Shoulder, NOS Arm <input type="checkbox"/> Arm, NOS Elbow <input type="checkbox"/> Elbow, NOS Forearm <input type="checkbox"/> Wrist <input type="checkbox"/> Forearm, NOS Hand <input type="checkbox"/> Metacarpal <input type="checkbox"/> Finger, NOS <input type="checkbox"/> Hand, NOS	Hip <input type="checkbox"/> Gr. trochanter <input type="checkbox"/> Femoral neck <input type="checkbox"/> Hip, NOS Leg (Upper) <input type="checkbox"/> Quadriceps <input type="checkbox"/> Hamstring <input type="checkbox"/> Femur <input type="checkbox"/> Upper leg, NOS Leg (Lower) <input type="checkbox"/> Tibia <input type="checkbox"/> Fibula <input type="checkbox"/> Gastrocnemius muscle <input type="checkbox"/> Lower leg, NOS Knee <input type="checkbox"/> Medial collateral ligament <input type="checkbox"/> Lateral collateral ligament <input type="checkbox"/> Anterior cruciate ligament <input type="checkbox"/> Posterior cruciate ligament <input type="checkbox"/> IT Band <input type="checkbox"/> Medial meniscus <input type="checkbox"/> Lateral meniscus <input type="checkbox"/> Other meniscus <input type="checkbox"/> Patella <input type="checkbox"/> Patella tendon <input type="checkbox"/> Patellofemoral joint <input type="checkbox"/> Tibial plateau <input type="checkbox"/> Knee, NOS Ankle <input type="checkbox"/> Achilles <input type="checkbox"/> Lateral ligament <input type="checkbox"/> Medial ligaments <input type="checkbox"/> Ankle, NOS Foot <input type="checkbox"/> Metatarsal <input type="checkbox"/> Pes planus <input type="checkbox"/> Pes cavus <input type="checkbox"/> Plantar fascia <input type="checkbox"/> Sessmoid <input type="checkbox"/> Toe, NOS <input type="checkbox"/> Foot, NOS	Back & Spine <input type="checkbox"/> C-spine area <input type="checkbox"/> T-spine area <input type="checkbox"/> L-spine area <input type="checkbox"/> Back or spine, NOS Rib <input type="checkbox"/> Rib, NOS 4. Injury Category (check one) <input type="checkbox"/> Overuse <input type="checkbox"/> Traumatic <input type="checkbox"/> Other <input type="checkbox"/> Unknown 5. Type of Injury (check one) <input type="checkbox"/> Normal exam <input type="checkbox"/> Abrasion/Laceration <input type="checkbox"/> Arthritis <input type="checkbox"/> Bursitis <input type="checkbox"/> Contusion <input type="checkbox"/> Dislocation <input type="checkbox"/> Fasciitis <input type="checkbox"/> Fracture <input type="checkbox"/> Ingrown toenail <input type="checkbox"/> Instability <input type="checkbox"/> Muscle spasm <input type="checkbox"/> Neuropathy <input type="checkbox"/> Osteochondral defect <input type="checkbox"/> Pain <input type="checkbox"/> Radiculopathy/Radiculitis <input type="checkbox"/> Rupture <input type="checkbox"/> Shin splints <input type="checkbox"/> Strain <input type="checkbox"/> Spondylosis <input type="checkbox"/> Sprain <input type="checkbox"/> Stress fracture <input type="checkbox"/> Stress reaction <input type="checkbox"/> Synovitis <input type="checkbox"/> Subluxation <input type="checkbox"/> Tendinitis <input type="checkbox"/> Tear <input type="checkbox"/> Other <div style="border: 1px solid black; width: 100px; height: 1.2em; display: inline-block;"></div> <input type="checkbox"/> Unknown
---	--	---

6. Type of Visit (check one) <input type="checkbox"/> Initial Visit For This Injury <input type="checkbox"/> Follow Up Visit For This Injury 7. Disposition (check one) <input type="checkbox"/> No profile <input type="checkbox"/> Profile <input type="checkbox"/> Quarters <input type="checkbox"/> Hospitalized <input type="checkbox"/> Other <input type="checkbox"/> Unknown 8. Number of Profile or Quarters Days <div style="border: 1px solid black; width: 40px; height: 1.2em; display: inline-block;"></div> DAYS <input type="checkbox"/> EPTS (recommended) <input type="checkbox"/> MEB 9. Consultation (check one) <input type="checkbox"/> None <input type="checkbox"/> Orthopedics <input type="checkbox"/> Podiatry <input type="checkbox"/> Physical Therapy <input type="checkbox"/> Gen surgery (cast clinic) <input type="checkbox"/> Other	Record only one injury (the most serious).
---	---